TAXONOMIC INDEX

Copeia 2013, Nos. 1–4

NEW NAMES

Acrobrycon
ortii 604–611
starnesi 604–611
Arapaima ***leptosoma*** 470–475
Caristius
digitus 425–432
meridionalis 426–431, 432
Characidium
amaila 454–463
papachibe 52–57
Cyphocharax ***aninha*** 627–633
Doumea ***skeltoni*** 577–581
Hemigrammocharax ***rubensteini*** 31–37
Hyphessobrycon ***kayabi*** 613–618
Hypomasticus ***lineomaculatus*** 464–468
Hypopygus ***benoneae*** 232–237
Incilius ***majordomus*** 9–10
Leporinus ***parvulus*** 241–244
Listrura ***depinnai*** 641–646
†*Lophionotus* ***chinleana*** 582–603
Microcaecilia ***savagei*** 223, 225–226, 228–229
Oreolalax ***sterlingae*** 214–221
Oryzias
asinua 403–414
wolasi 403–414
Parotocinclus ***jequi*** 435–439
Platyberyx
paucus 417–423, 432
pietschi 417–424, 432
rhyton 417–424, 432
Tatia
carolae 396–399, 401
marthae 398–401
Trachycephalus ***helioi*** 634–638
Typhlobelus ***auriculatus*** 441–453

NEW AND ESTABLISHED NAMES

Abramidina 14, 19
Abramidinae 14
Abramites 244
hypsenolotus 244
Acanthaster 347
Acanthoscyllium sahelalmae 379
Acantopsis choirorhynchos 16, 19
Acarina 105
Acer 249
rubrum 255–256
saccharum 677
Acestrocephalus 35
Acheilognathinae 16
Acheilognathus typus 16, 19
Achillea millefolium 65
Achylasp 98
Acris crepitans 562–566
Acrobrycon 604–611
ipanquianus 604–611
ortii 604–611
starnesi 604–611
tarijae 604–611
Acrodytes
inflata 639
modesta 639
Acropoma 25
japonicum 27
Acropomatidae 23–30
Actinemys marmorata 142–153
Adinia xenica 91
Adrianichthyidae 403–414
Adrianichthys 413
kruyti 405
oophorus 405, 413
poptae 405
roseni 405
Afrolimon pergrinum 194
Agalychnis 163–164
callidryas 160, 163
Agropyron 65
Akistrodon piscivorus 489
Akmoniston 385
Alaria alata 105, 107
Alburniformes 14, 19
Alburninae 19
Alburnus 14
alburnus 19, 106
Alestidae 35, 617
Alligator 518
mississippiensis 268–276, 517–521
sinensis 518–519, 522–523
Alligatoridae 518
Alopias 386
Amblypharyngodon mola 16, 19
Amblytrypauchen arctocephalus 29
Ambystoma 758
californiense 786
laterale-jeffersonianum 254–261
maculatum 284–291, 338–345, 514
mavortium 786
opacum 285
talpoideum 284–291
tigrinum 786
Amia 24, 587–590, 600
(Amioides) 23
calva 586
grossidens 26
Amioides 23–30
grossidens 23–30
polyacanthus 23–30

© 2013 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/OT-13-138

Ammocryptocharax 55
elegans 462
vintonae 462
Amolops
afghanus 798
cucae 220
marmoratus 798
minutus 220
splendissimus 220
Amphibia 223–231
Amphiliidae 577–581
Anas platyrhynchos 104
Anax junius 691–698
Anaxyrus
americanus 251
canorus 8
fowleri 647–652
terrestris 543–551
Ancherythroculter 15
Aneides
ferreus 785
vagrans 785
Anolis 201, 203–204
conspersus 204
garmani 204
grahami 201, 203–204
jubar 201–205
j. albertschwartzi 201–205
j. maislensis 201–205
j. oriens 201–205
opalinus 204
valencienni 204
Anostomidae 464–469
Antennarius 663
commerson 664
multiocellatus 664–665
subtere 29
Anura 213–222, 634–640, 798
Apareiodon agmatos 462
Aphyocharacidium 35
Aphyocypris 13–22
chinensis 17, 19
Aphyoditeinae 35
Aplodinotus grunniens 740–749
Apogonidae 23–30
Araneae 105
Aranei 105
Arapaima 38–51, 470–476
agassizii 38–51, 470–476
arapaima 38–40, 42, 44, 470–476
gigas 38–51, 470–476
leptosoma 470–475
mapae 38–42, 44, 46, 48, 470–476
sp. 39–40, 42, 44–45, 48, 471, 473
†*Araripelepidotes* 582, 587–590, 600
temnurus 599
Ardea cinerea 104
Argenteohyla altamazonica 639
Ariomma parini 772
Ariommatidae 772
Arnoglossus tapeinosoma 29
Artemia 121
Aspidoparia 14
Aspidoscelis
arizonae 366–377
inornata 366–377
i. arizonae 366–377
i. gypsi 372–372
i. heptagramma 367
i. junipera 367
i. llanurus 366–377
pai 366–377
uniparens 367, 369–371, 374
velox 371
Astiotrema monticelli 107
Astroblepidae 462
Astyanax
ipanquianus 605
scabripinnis 617
Asymbllepharus himalayanus 178
Atelopus 8, 11
Atherinomorpha 403–414
Atherinops affinis 731
Atherinopsis californiensis 731
Atlantis 622
Atopichthys phillipsi 190
Atractosteus 587–589, 600
Auchenipteridae 396–402

Balitoridae 16
Barbatula toni 16, 19
Barbini 13
Barbonymus gonionotus 16, 19
Barbus
barbus 16, 19
trimaculatus 16, 19
Barilius 14
vagra 16, 19
Barisia imbricata ciliaris 714
Barycholos 760
Bassiana duperreyi 132
Bathylagichthys parini 772
Bathylagidae 772
Bathypteroidae 772
Bathypterois parini 772
Batrachochytrium dendrobatidis 97
Batrachoseps 786
attenuatus 786
catalinae 364
diabolicus 786
incognitus 786
kawia 786
leucopus 364
luciae 786
major 364
minor 786
regius 786
Belonesox belizanus 91
Beloniformes 403–414
Belostoma flumineum 691–698
Beryx 418
Biston betularia 782
Bivalvia 105
Bivibranchia velox 449
Blarina 806
Boidae 179
Boiga irregularis 100
Bolitoglossa savagei 760
Bombina bombina 702

Borophryne apogon 669
Borrelia burgdorferi 796
Bothrops atrox 684–690
Botiidae 16
Brachionichthys hirsutus 787
Brachycera 105
Bufo 8
  alvarius 363
  californicus 786
  debilis 760
  fastidiosus 8, 11
  fowleri 647–652
  guttatus 8
  holdridgei 8
  japonicus 335
  marinus 8, 335, 539, 548
  mauritanicus 335
  melanostictus 332–337
  peripatetes 8–9
  valliceps 8
  variegates 8
  viridis 706
Bufonidae 8–12, 706
Bungarus caeruleus 179
Buprestidae 105
Buteo buteo 69

Cabdio 14
  morar 16, 19
Cacophis squamolosus 164
Caecilia
  gracilis 229
  iwokramae 229
  pressula 229
  tentaculata 229
Caeciliidae 229
Caiman 518, 521
  crocodilus 275, 519, 521
  crocodylus 520
  latirostris 269, 275, 518–521
  yacare 518–521
Callanthidae 772
Callanthias parini 772
†*Callipurbeckia*
  minor 587–589, 600
  notopterus 599
Campostoma 354
  anomalum 353–354
  obligolepis 354
  pullum 354
Candidia 13–22
  barbatus 17, 19
Carabidae 105
Carapa guianensis 224
Carapidae 772
Carassius auratus 16, 19, 277–283
Carcharhiniformes 379
Carcharhinus leucas 378
Carcharias 386
Carcharodon 378–379, 383, 385–386
  carcharias 351
Caretta caretta 307, 309
Carettochelys insculpta 163
Carex sp. 104
Caristiidae 415–434
Caristius 415–434
  andriashevi 415, 417–418
  digitus 425–432
  fasciatus 426–432
  groenlandicus 415, 428
  japonicus 415, 417, 424–425, 429–430
  macropus 425–432
  meridionalis 426–431, 432
  opalescens 421
  sp. 432
Carpiodes velifer 354
Catastomidae 354
Catostomidae 16, 390–395
Catostominae 16
Catostomus 390
  catostomus 121, 390
  commersoni 121, 390
  commersonii 16, 19
Caulophrynidae 668
Ceanothus sp. 147
Centaurea solstitialis 65
Centrarchidae 354
Centrolenella savagei 760
Centromochlinae 396–402
Centromochlus musaicus 401
Centrophryne spinulosa 666–669
Centrophrynidae 666–669
Cepea nemoralis 782
Ceratias 668
Ceratiidae 668–669
Ceratioidei 666, 668
Ceratophyllum demersum 103–104
Ceratopogonidae 105
Cetichthys parini 772
Cetomimidae 772
Cetopsis 610
Cetorhinus 789
Chalceus macrolepidotus 315
Chaenobryttus gulosus 364
Chaetodontidae 798
Chanodichthys 15
Chanos chanos 14, 16, 19, 406–407, 413
Chara 81
Characidae 35, 604–621
Characidiinae 55, 454
Characidium 35, 52–57
  aff. *zebra* 56
  alipioi 455–456, 461
  amaila 454–463
  bahiense 56
  cf. *pteroides* 56
  boavistae 454, 456–458, 462
  bolivianum 454, 456–457, 461, 462
  cf. *boavistae* 458, 462
  cf. *bolivianus* 458
  crandellii 56, 456
  declivirostre 56, 455–456, 462
  etheostoma 56
  fasciatum 454–459, 461
  gomesi 454–456, 458, 461–462
  grajahuensis 455–456, 461
  hasemani 56, 462
  heinianum 52–53, 55
  heirmostigmata 52–57

japuhybense 455–456, 461
lauroi 455–456, 461
longum 53, 55–56
nupelia 55–56
occidentale 55
orientale 55
papachibe 52–57
pellucidum 53, 55–56
pteroides 53, 55–56
purpuratum 454, 456–458, 461–462
rachovii 55
serrano 52–57
stigmosum 55–56
timbuiense 455–456, 461
vestigipinne 55
vidali 455–456, 461
xavante 55–56
zebra 455, 463
Characiformes 16, 31–37, 52–57, 312–323, 454–469, 604–621, 627–633, 800
Characinae 35
Charax 35
Chaunus arenarum 563
Chedrini 14
Cheilodipterus 23, 28
polyacanthus 23, 26, 29
singapurensis 29
Cheirodontidae 618
Cheirodontinae 35
Chela 13–15
cachius 16, 19
Chelae 13
Chelaethiops bibie 16, 19
Chelodina rugosa 117
Chelonia mydas 307
Chelydra serpentina 58–61
Chiasmodontidae 772
Chilomeniscus savagei 760
Chilophryne conifera 8
Chiloscyllium 379
Chironomidae 105, 616
Christinus marmoratus 132
Chromobotia macracantha 16
Chrosomus 750, 752
cumberlandensis 755
erythrogaster 755
saylori 750
sp. cf. *saylori* 750–756
tennesseensis 755
Chrysemys
picta
p. bellii 570
p. marginata 570
Chrysomelidae 104–105
Chrysosporium 97–102
Cichlasoma urophthalmus 80–96
Cichlidae 449, 462
Circaetus gallicus 69, 140
Cladocera 105, 616
Clarias
depressus 363
gariepinus 364
sp. 410
Clinostomus funduloides 364, 625
Clupeidae 354
Cnemidophorus
arizonae 367
inornata 367
i. junipera 367
inornatus 367
Cobitidae 16
Cobitis striata 16, 19
Cobitoidea 15–16
Coccinelidae 105
Cochranella savagei 760
Codonocephalus urnigerus 105–106
Coleoptera 104–105, 754
Coluber
rhodorhachis 179
ventromaculatus 179
Colubridae 132–141, 179
Conger oceanicus 190
Congoglanis 580
Congridae 191, 772
Coniophanes fissidens 132, 139–140
Contia
longicaudae 785
tenuis 785
Copepoda 105
Copionodontinae 444, 451
Coranthus 23–30
polyacanthus 23–30
Corbicula 478–481
fluminea 477–478
sp. 477–484
Coris aygula 787
Corixidae 105
Cosmocerca ornata 106
Cosmopolitodus hastalis 387
Cottidae 354
Cottus
caeruleomentum 624–625
cf. *bairdii* 355
Cranophryne 8
Cranopsis 8
fastidiosus 8
Creagrutus muelleri 239
Cremnobates 363
Crenuchidae 35, 52–57, 454–463
Crepidophryne 8, 11
chompipe 8
epiotica 8
guanacaste 8
Crocodylia 517–526
Crocodylidae 523
Crocodylus 523
acutus 275, 517–520, 522–523
crocodilus 518, 524
intermedius 518–520, 522–523
johnstoni 520, 522
mindorensis 521–523
moreletii 518, 521–523
niloticus 517–523
novaeguineae 269, 519–520, 522–523
palustris 517, 519–522
porosus 275, 518–523
rhombifer 519, 521–523
siamensis 519–520, 522–523

suchus 519–520, 522
Crotalidae 707
Crotalus 490, 707
adamanteus 489
atrox 707, 786
durissus 489
lepidus 708
l. klauberi 714–715
l. lepidus 714–715
l. morulus 707–716
mitchellii 785
oreganus 486
o. oreganus 485–492
pricei 485, 490
scutulatus 490, 786
s. scutulatus 707
simus 707
stephensi 785
tigris 490
unicolor 489
viridis 486
Crotaphytus
collaris 190
vestigium 786
Cryptobranchus alleganiensis 514
Cryptopsaras couesii 669
Cryptotomus 363
Ctenoluciidae 617
Ctenopharyngodon 13–22
idella 17, 19
Culter 13–22
alburnus 15, 17, 19
Cultrops 14
Curculionidae 104–105
Curimatella 630
dorsalis 632
immaculata 632
Curimatidae 627–633
Curimatopsis 631
Cyclanthaceae 685
Cycleptinae 16
Cycleptus elongatus 16, 19, 319
Cynopoecilus 644
Cyphocharax 627–633
abramoides 627–628
aninha 627–633
aspilos 627–628
biocellatus 628–630
derhami 628
festivus 628, 632
gangamon 627–628, 630–632
gilbert 628, 630
gillii 628, 630
gouldingi 628, 630, 632
helleri 628–630, 632
laticlavius 628–629
leucostictus 627–628
magdalenae 628
meniscaprorus 628, 630–631
mestomyllon 628, 630–632
microcephalus 628
modestus 628, 630
multilineatus 628–629
nagelii 628
nigripinnis 628, 631–632
notatus 628–629
oenas 628, 630–631
pantostictos 628–630
piluropisis 629
pinnilepis 627–629
platanus 628
plumbeus 628, 632
punctatus 627–631
saladensis 628, 630–631
santacatarinae 628–630
signatus 627–632
sp. 632
spilotus 629–630, 632
spiluropsis 630, 632
spilurus 629–630, 632
stilbolepis 628, 632
vanderi 627, 629–631
vexillapinnus 627–632
voga 629–630
Cyprinella
cf. *spiloptera* 354–355
lutrensis 17, 19
spiloptera 354
whipplei 354
Cyprinidae 13–22, 354, 740, 800
Cypriniformes 13–22
Cyprinodon variegatus 80–96
Cyprinodontiformes 527–538
Cyprinoidea 15–16
Cyprinus carpio 16, 19, 281, 353

Daboia russelii 178
Dalatiidae 772
Danio 14
dangila 17, 19
rerio 17, 19, 528
Danionella dracula 17, 19
Danioninae 13–16, 19, 800
Danionini 14
†*Dapedium* 587–589, 600
pholidotus 599
punctatus 599
Daphnia lumholtzi 300
Darevskia 181
Deirochelys reticularia 149
Demansia psammophis 164
Dendroaspis polylepis 198
Dendrophidion dendrophis 132–141
Dendrophryniscus 11
†*Dentilepisosteus* 587–589, 600
Derhamia hoffmannorum 462
Dermochelys coriacea 309
Desmognathus 262–267, 499–506
aeneus 262–267, 499–500
conanti 262
folkertsi 499–506
fuscus 262, 363, 503
marmoratus 499
monticola 262–267, 499–500
ocoee 262–267, 499–500, 512–516
orestes 262
organi 262
quadramaculatus 262–267, 499–506
santeetlah 262

wrighti 262–267
Deuterodon 618
langei 618
Devario auropurpureus 17, 19
Diaphus parini 772
Diapoma 604–605, 610
Diapomini 604–605, 610
Dinolestidae 24
Dionda diaboli 73
Diplodiscus subclavatus 104, 106
Diptera 105, 754
Diretmichthys parini 772
Diretmidae 772
Distichodontidae 31–37, 800
Distoechodon 15, 19
Dolichopteryx parini 772
Dolomedes
fimbriatus 104
plantarius 104
Donax variabilis 782
Dorosoma petenense 353
Doumea 577–581
angolensis 580–581
chappuisi 578, 580–581
gracila 580–581
reidi 580–581
sanaga 578, 580–581
skeltoni 577–581
thysi 578, 580–581
stilicauda 581
typica 580–581
Doumeinae 577–581
Dreissena 481
bugegensis 481
polymorpha 481
sp. 477–484
Drosophila melanogaster 678
Duttaphrynus melanostictus 332

Echinochloa pyramidalis 304
Eirenis persicus 181
Elachistocleis bicolor 706
Elachocharax 55
Elaphe 168
obsoleta 507
Elapidae 179
Eleocharis cellulosa 82
Elephenor 424–425, 430
macropus 429
Elephenoridae 425, 430
Eleutherodactylus
adamastus 760
coqui 335
jota 760
Elimia
potosiensis 481, 483
virginica 478
Elliptio complanata 478
Elopichthys 15, 19
Emydoidea blandingii 307, 571
Emys orbicularis 308
Encyclometra colubrimurorum 105–107
Engraulis mordax 731
Enulius 132, 139
Ephemeroptera 105
Epicrionops 226, 228, 230
bicolor 229
lattivittatus 229
marmoratus 229
niger 223–231
nigrus 224
parkeri 229
peruvianus 229
petersi 229
sp. 229
Epigonidae 24, 27, 772
Epigonus 24
pandionis 27
parini 772
Ericaulaceae 460
Erythrinidae 312–323, 617
Erythroculter 15
Eulamprus heatwolei 541
Esomus danricus 17, 19
Esox lucius 107
Etheostoma 154–159
asprigene 354–355
barrenense 155, 157
caeruleum 155–156
camurum 353–354
(Catonotus) 155
cf. *aspringene* 354–355
cf. *zonale* 355
chlorosoma 353
flabellare 353, 355, 656
fusiforme 624
histrio 355
hopkinsi 154–159
h. binotatum 155
juliae 658
luteovinctum 154–159
maculatum 353
(Nanostoma) 155
neopterum 656
nigripinne 155–157
okaloosae 653–659
(Oligocephalus) 154–156
olmstedi 625
podostemone 656
spectabile 156
(Ulocentra) 155
variatum 355
zonale 155, 157, 355
zonatum 355
Eugomphodus 379
Eumeces
brevirostris pineus 714
taeniolatus 179
Euphorbia
mauritanica 194
seguieriana 65
Eurycea bislineata 363, 512
Eurylepis 179
Eustomias parini 772
Euterpe oleracea 224
Eutropis 179, 800
Exocoetidae 768

Fagus grandifolia 677
Floridichthys carpio 91

Fundulidae 354, 527–538
Fundulus
chrysotus 91
confluentus 80–96
grandis 91
heteroclitus 527–538
h. heteroclitus 527
h. macrolepidotus 527
similis 91

Gaidropsarus parini 772
Galeocharax 35
Gambelia copeii 785
Gambusia 72–79, 248
affinis 73–75, 77–78, 91, 156, 248–253
cf. *clarkhubbsi* 72–79
clarkhubbsi 72–79
geiseri 75, 156
heterochir 73
holbrooki 74, 80–96, 251
krumholzi 72–79
nobilis 73, 78
speciosa 72–79
Gasterosteus 156
aculeatus 127
Gastrophryne carolinensis 543–551
Gastropoda 105
Gavialidae 523
Gavialis gangeticus 517–520, 523–524
Gekkonidae 132
Gekkota 206
Geochelone pardalis 570
Gerridae 104–105
Geryichthys 55–56
sterbai 52–53, 55
Gigantactis savagei 760
Ginglymodi 582
Glanapteryginae 444, 641–646
Glanapteryx 441, 443, 641–642, 645
anguilla 441, 452, 645
niobium 452
Glyptemys
insculpta 329
muhlenbergii 324–331, 724
Gnathophis parini 772
Gnathopogon elongatus 17, 19
Gobio gobio 17
Gobioninae 17
Gonorhynchiformes 16
Gonorynchus greyi 16, 19
Gopherus agassizii 571
Gorgodera
asiatica 106
pagenstecheri 106
varsoviensis 106
Grandisonia brevis 228
Graptemys 477, 483, 571, 723
caglei 723–728
flavimaculata 723
geographica 477–484, 723–724, 727
kohnii 723, 727
Gymnetrus hawkinii 356
Gymnocypris przewalskii 16, 19
Gymnophiona 223–231
Gymnorhamphichthys sp. 449
Gymnothorax
parini 772
punctarenae 190, 363
Gymnotiformes 232–237
Gymnotus 610
aff. *pantherinus* 644
Gyrinocheilidae 16
Gyrinocheilus aymonieri 16, 19

Halecomorphi 582
Hamamelis virginiana 255–256
Haplophryne mollis 669
Hemibarbus barbus 17, 19
Hemibrycon
ipanguianus 605
ipanquianus 605
†*Hemicalypterus weiri* 599
Hemiculter 15
Hemiculterella 15
Hemigrammocharax 31–37
angolensis 31, 35
lineostriatus 32, 35
machadoi 31, 35
minutus 31, 35
monardi 31
multifasciatus 31, 35
ocellicauda 31, 35
rubensteini 31–37
uniocellatus 31, 35
wittei 31, 35–36
Hemigrammocypris 13–22
rasborella 17, 19
Hemigrammus 612
sp. 616
Hemiodontidae 449
Hemiphractidae 180
Hemiptera 104–105, 616
Hemiscyllium 379
Heterandria formosa 91
Heterodon platirhinos 363, 507–511
Heterotis 47
Hiodon tergisus 353
Histrio histrio 782
Hologymnosus annulatus 787
Homaloptera leonardi 16, 19
Homo sapiens 179
Hoplias 313
cf. *malabaricus* 618
malabaricus 313, 315–316, 318–320
microlepis 312–323
Hoplodactylus duvaucellii 781
Horadandia atukorali 17, 19
Howella parini 772
Howellidae 24, 772
Hybognathus 353
Hydrobiidae 477–484
Hydromantes platycephalus 512
Hydrophiidae 179
Hyla
adenoderma 639
arborea 702–703
chrysoscelis 796
cinerea 543–551, 796
hadroceps 634

japonica 706
lichenosa 639
macrotis 639
nana 706
nigropunctata 639
palpebrogranulata 639
raniceps 706
salvaje 760
squirella 543–551
versicolor 562
wettsteini 639
zernyi 639
Hylidae 634–640, 706
Hymenoptera 105, 754
Hypentelium nigricans 16, 19, 121
Hyperoliidae 781
Hyperolius cinnamomeoventris 164
Hyphessobrycon 612–621
aff. *stegemanni* 619
agulha 618
amandae 618
anisitsi 617–618
arua 618
balbus 616–618
bentosi 618
bifasciatus 612–621
bleheri 618
cf. *copelandi* 618
cf. *malabaricus* 618
coelestinus 618
cyanotaenia 612–621
duragenys 616–618
elachys 35, 617–618
eques 618
flammeus 618
griemi 618
hamatus 616–618
heliacus 612–621
herbertaxelrodi 618
heterorhabdus 618
hexastichos 618
igneus 612–621
kayabi 613–618
langeanii 616–617, 629
loweae 612–621
luetkenii 619
megalopterus 619
melanostichos 619
meridionalis 618
micropterus 619
moniliger 617, 619
nicolasi 618
notidanos 617
peugeoti 612–621
pulchripinnis 619
reticulatus 616–617, 619
rutiliflavidus 617
santae 616–617, 619
savagei 760
socolofi 617
stegemanni 618
togoi 618
vilmae 619
vinaceus 616, 619
Hypomasticus 464–469
despaxi 464–469
garmani 464
julii 464
lineomaculatus 464–468
megalepis 464, 467
mormyrops 464
pachycheilus 464
thayeri 464
Hypophthalmichthyina 19
Hypophthalmichthyinae 13–22
Hypophthalmichthys 13–22
molitrix 15, 18–19
nobilis 18–19
Hypopomidae 232–237
Hypoptopomatinae 435–440
Hypopygus 232–237
benoneae 232–237
cryptogenes 232–233, 236
hoedemani 232–237
isbruckeri 232, 234, 237
lepturus 232–237
minissimus 232–237
neblinae 232, 234, 236
nijsseni 232, 234, 237
ortegai 234, 236
Hypsopanchax 363

Ichthyococcus parini 772
Icosiella neglecta 106
Ictaluridae 354
Ictalurus
furcatus 353
punctatus 15, 16, 19, 277–283
Idiocranium russelli 228
Imparfinis pristos 449
Incilius 8–12, 363
bocourti 11
chompipe 11
coccifer 11
coniferus 8, 11
(Crepidophryne) 11
epiotica 11
fastidiosus 8, 10–11
guanacaste 11
holdridgei 10–11
ibarrai 11
karenlipsae 8
majordomus 9–10
nebulifer 562–566
periglenes 8
peripatetes 8–12
†*Isanichthys* 587–589, 600
Ischikauia 13–22
steenackeri 18–19
Isopoda 105
Isthmiophora melis 104, 106
Isurus 379, 383, 386
oxyrinchus 364, 385
oxyrinhus 347
punctatus 378
tigris 364
Ixodida 754

Jaera sarsi 104
Jaydia striata 29
Jordanella floridae 91
Juncus effusus 657
Jupiaba acanthogaster 618

Kinosternon
  durangoense 571
  sonoriense 571
Klausewitzia 55
†*Kyphosichthys grandei* 599

Labeo senegalensis 16, 19
Labeonini 13
Labidesthes sicculus 292–302
Labridae 493–498
Lacerta agilis 103–104
Laemolyta 244
Lamna 378–389
  cornubica 378
  ditropis 378–389
  nasus 378–389
  philippii 378
  whitleyi 378
Lamnidae 378–389
Lamniformes 379, 385
Lampridiformes 356
Lampropeltis getula 786
Lampropholis delicata 160–165
Laubuca 13
  laubuca 17, 19
Laubucae 13
Laudakia agrorensis 178
Lebiasina
  aff. *bimaculata* 619
  ardilai 462
Lebiasinidae 462, 617
Lefua echigonia 16, 19
Leiolopisma schmidti 364
Leiuperidae 706, 798
Lemna trisulka 104
Lepidarchus 35
Lepidochelys olivacea 117
Lepidoptera 105
†*Lepidotes* 582
  elvensis 599
  gigas 588–590, 599–600
  microrhis 599
  semiserratus 588–590, 600
  sp. 599
Lepisosteiformes 582
Lepisosteus 587–589, 600
Lepomis 85–86, 88, 91, 93
  gibbosus 364
  gulosus 88, 364
  macrochirus 88, 251, 277–283
  marginatus 91
  microlophus 88
  punctatus 91
Leporellus 244, 464–465
Leporinus 238–247, 464–465, 467
  agassizi 241
  agassizii 238–239, 242, 245
  amae 465–466
  amblyrhynchus 465–466
  arcus 465–467
  bistratus 465–466
  britskii 465–466
  cf. *moralesi* 242
  geminis 465–466
  jamesi 244–245
  (Leporinops) 239, 244
  margaritaceus 245
  melanopleura 465–466
  moralesi 238–247
  muelleri 238–247
  mulleri 239
  mülleri 239, 241
  nigrotaeniatus 238–239, 245
  parvulus 241–244
  punctatus 242
  reinhardti 244–245
  semivittatus 245
  sexstriatus 467
  striatus 242, 244–245, 465–467
  taeniatus 242, 245, 465–466
  taeniofasciatus 465–466
  uatumaensis 238–247
  unitaeniatus 465–466
  vanzoi 244, 465–466
Leptagoniates pi 35
Leptobarbinae 17
Leptobarbus hoevenii 17, 19
Leptobotia mantschurica 16, 19
Leptobrachium
  boringiae 213–215
  echinatum 220
  xanthospilum 213–215
Leptocharacidium 55
Leptocypris niloticus 17, 19
Leptodactylidae 706
Leptodactylus
  chaquensis 706
  fuscus 706
  podicipinus 706
  savagei 760
Leptolalax pluvialis 220
†*Leptolepis* 587–589, 600
  coryphaenoides 586
Leptophallus nigrovenosus 107
Leptotyphlops macrorhynchus 179
Leuciscinae 17, 19
Leuciscus
  oxygaster 14
  vandoisulus 364
Leuresthes
  sardina 730, 736
  tenuis 729–739
Liasis fuscus 64
Linophryne parini 772
Linophrynidae 668–669, 772
Liolaemidae 206–212
Liolaemus 209
  chiliensis 206–212
Liparidae 772
Liriodendron tulipifera 677
Listrura 441, 443, 641–646
  boticario 641–646
  camposi 452, 641–645

costai 641–645
depinnai 641–646
nematopteryx 452, 641–645
picinguabae 452, 641–645
sp. 644–645
tetraradiata 452, 641–645
Lithobates
areolatus 552–561
a. areolatus 559
a. circulosus 559
berlandieri 562–566
capito 552
catesbeianus 543–551, 691–698
clamitans 248–253, 691–698
(Rana) 691–698
sevosus 552
sphenocephalus 543–552
sylvaticus 251, 553, 562, 670, 796
Lithogenes villosus 461
Litoria
aurea 562
ewingii 562
Longiculter 14–15
Lontra canadensis 61, 150
Lophelia pertusa 660
Lophiidae 660–665
Lophiiformes 660–669
Lophiodes
beroe 663
caulinaris 663
kempi 663
reticulatus 663
sp. 663
†*Lophionotus* 582–603
chinleana 582–603
sanjuanensis 582–603
Lophogobius cyprinoides 91
Loricariidae 435–440, 462
Lota lota 107
Lotidae 772
Lucania
goodei 91
parva 91
Luciobrama 19
Luciosoma 14
setigerum 17, 19
†*Luoxiongichthys* 587–589, 600
Luxilus 354
chrysocephalus 354
cornutus 354
Lygaedae 105
Lythrurus 354
cf. *umbratilis* 355
fasciolaris 353
fumeus 354

Mabuya dissimilis 179
Macrhybopsis storeriana 354
Macrochelys temminckii 58–63
Macrochirichthys 13–22
macrochirus 18–19
Macrodera longicollis 105, 107
fegerti 587–589, 600
lennieri 587–589, 600
†*Macrosemius* 587–589, 600
Macrouridae 771–772
Malacanthus latovittatus 787
Malayochela maassi 17, 19
Manzanita sp. 147
Marantaceae 685
†*Masillosteus* 587–589, 600
Masticophis
bilineatus 786
flagellum 198, 785
fuliginosus 785
taeniatus 786
Mastigodryas bifossatus 132–141
Mazarunia 462
Mecistops cataphractus 519–520, 522
Medicago coerulea 65
Megachasma 379
Megalobrama 13–22
amblycephala 18–19
Megaloptera 754
Megophryidae 213–222
Megophrys omeimontis 213–215
weigoldi 213
Megaselachus megalodon 387
Melamphaes parini 772
Melamphaidae 772
Melanocetidae 668
Melanocetus johnsonii 668
Melanocharacidium 55
Melanophryniscus 11
Melanosuchus niger 518–521
Menidia 292
audens 292–302
beryllina 292, 353
menidia 735
peninsulae 80–96
Mesobola brevianalis 15, 17, 19
Metzia 13–22
formosae 18–19
lineata 18–19
Microcaecilia 223–231
albiceps 223, 226, 229
grandis 223, 226, 228–229
iyob 223, 226, 228–229
rabei 223, 226, 228–229
rochai 223, 226, 228–229
savagei 223, 225–226, 228–229, 760
sp. 1 224
sp. 2 224, 228
supernumeraria 228–229
taylori 223, 226, 228–229
trombetas 223, 226, 228–229
unicolor 223, 226, 228–229
Microcambeva
barbata 645
ribeirae 645
Microcharacidium 55
Microdevario kubotai 17, 19
Microgobius gulosus 80–96
Microhylidae 706, 781
Micropterus salmoides 277–283, 654
Micropanchax 363
Microrasbora rubescens 17, 19
Microschemobrycon 35
Microtus 806

sp. 69
Milvus migrans 104
Mimosa pigra 304
Misgurnus fossilis 107
Mitsukurina 379
Mnierpes macrocephalus 348
Moenkhausia
ceros 616
cf. *lepidura* 616
collettii 616
cotinho 616
oligolepis 616
Mollisquama parini 772
Montivipera 64
raddei 64–71
Mora excelsa 224
Morelia spilota imbricata 69
Moridae 772
Morone
americana 353
chrysops 353
Moxostoma 125–126, 390–395
anisurum 354
carinatum 121–126, 390, 392
duquesnei 121–126, 390–395
erythrurum 121–126, 390–395
hubbsi 121–126
macrolepidotum 121–126
melanops 121–126
valenciennesi 121–126, 354
Mugil cephalus 353
Mullidae 191
Mullus phillipsi 191
Muntiacus 521
Muraenidae 772
Mycteroperca venenosa 798
Myctophidae 772
Mylogpharyngodon 19
Myloplus planquettei 315
Myotis 806
Myriopholis 179
Myxocyprinus asiaticus 16, 19

Naja oxiana 179
Nannaethiops 35
Nannizziopsis viresii 97
Nannocharax 31–37
ansorgii 32, 35–36
brevis 35–36
elongatus 35–36
fasciatus 35–36
intermedius 35–36
latifasciatus 32, 35–36
lineomaculatus 32, 35–36
macropterus 35–36
maulicauda 35–36
niloticus 35–36
occidentalis 35–36
ocellicauda 31
parvus 32, 35–36
procatopus 35–36
pteron 35–36
reidi 35–36
rubrolabiatus 35–36
schoutedeni 35–36
seyboldi 35–36
taenia 35–36
usongo 32, 35–36
Nannophryne 11
Nannostomus beckfordi 619
Nanochloropsis 121
Natriceteres 139
Natrix 103
natrix 69, 103–110
stolata 100
tessellata 103–110
Neblinichthys 462
Nemacheilidae 16
Nematabramis 14
steindachneri 17, 19
Nematoda 106
Neobolini 14
Neobola bottegoi 17, 19
Neocaristius 415–416
Neoceratiidae 668
Neodiplostomum spathoides 107
Neolebias 35
Neopterygii 582–603
†*Neosemionotus* 587–589, 600
punctatus 599
Nepidae 105
Nerodia sipedon sipedon 139–140
Nezumia parini 772
Nicholsicypris 13–22
normalis 18–19
Nicholsina 190, 363
Nipponocypris 13–22
sieboldii 18–19
temminckii 18–19
Nippotaenia mogurndae 106
Nomorhamphus sp. 410
Norops 201
(Anolis) 201, 204
humilis 204
sagrei 201, 204
†*Notagogus* 587–589, 600
Notemigonus crysoleucas 17, 19
Notophthalmus viridescens 691–698
Notropis 354
amoenus 625
amnis 354–355
anogenus 353–354
ariommus 354
atherinoides 17, 19
buchanani 353–354
heterodon 354
heterolepis 354
percobromus 354
rubellus 354
shumardi 353–354
texanus 354
Noturus
gyrinus 282
nocturnus 353
stigmosus 354
Nudiantennarius subtere 29

†*Obaichthys* 587–589, 600
Ochetobius 13–22
elongatus 18–19

Odonata 105
Odontocharicidium 55
Oedemeridae 105
Oedipina savagei 760
Ollotis 8
  coerulescens 8
Oncorhynchus mykiss 354
Oneirodidae 668
Ophiohagus hannah 100
Ophiotaenia europaea 103–110
Ophisops jerdonii 178
Opisthioglyphe ranae 104, 106–108
Opisthoproctidae 772
Opsaridium peringueyi 17, 19
Opsariichthyinae 13–22
Opsariichthyini 19
Opsariichthys 13–22
  bendelisis 17, 19
  bidens 18–19
  pachycephalus 18–19
  uncirostris 18–19
Opsarius 14
  bendelisis 17, 19
Opsopoeodus emiliae 353
Orectolobus 379
Oreochromis sp. 410
Oreolalax 213–222
  chuanbeiensis 214–217
  granulosus 213, 216–217
  liangbeiensis 214–217, 221
  jingdongensis 214–217, 220
  lichuanensis 214–217
  major 214–217
  multipunctatus 214–217, 220
  nanjiangensis 214–217
  omeimontis 214–217
  pingii 214–217
  popei 214–217
  puxiongensis 213, 216–217, 220
  rhodostigmatus 214–217
  rugosus 214–217, 221
  schmidti 214–217
  sp. 214–217
  sterlingae 214–221
  weigoldi 213, 216–217
  xiangchengensis 214–217
Oreophrynella 11
Orthoptera 105
Oryzias 403–414
  asinua 403–414
  bonneorum 405, 413
  celebensis 405–407, 413
  eversi 405
  hadiatyae 405
  javanicus 405–407, 413
  latipes 413, 528
  marmoratus 405
  matanensis 405
  nebulosus 405
  nigrimas 405
  orthognathus 405
  profundicola 405
  sarasinorum 405, 413
  sp. 411–412
  wolasi 403–414
  woworae 403–414
Oscaecilia zweifeli 229
Osornophryne 11
Ostariophysi 604–611
Osteobrama 13
Osteoglossidae 38–51, 470–476
Osteoglossomorpha 38–51, 470–476
Osteolaemus 517, 520, 524
  tetraspis 518–519, 521–524
Osteopilus 543–544
  septentrionalis 543–551
Ostorhinchus
  atrogaster 29
  semilineatus 29
Ostracoda 105
Oswaldocruzia filiformis 106
Oxygaster 13–22
  anomalura 13, 18–20
  pointoni 13
Oxygastri 13–22
Oxygastrinae 13–22
Oxygastrini 14
Oxyporhamphidae 768
Oxytrichidae 98

Palaeoxantusia fera 760
Paleosuchus 518, 521
  palpebrosus 518–520, 524
  trigonatus 518–520
Pantherophis obsoletus 64, 68–69
Parabramis 15
Paracaristius 415–416, 432
  heemstrai 415
  maderensis 415, 421, 430
Paracheirodon axelrodi 35
Parachela 13–22
  maculicauda 18–19
  oxygastroides 18–19
  siamensis 18–19
  williaminae 18–19
Paraclinus 363
  fasciatus 363
Paralaubuca 13–22
  typus 18–19
†*Paralepidotus* 582, 587–590, 600
Paralepoderma cloacicola 106–107
Pararasbora 13–22
  moltrechti 18–19
Paratodus benedenii 387
Parazacco 13–22
  spilurus 18–19
Parodontidae 462
Parotocinclus 435–440
  arandai 436
  bahiensis 436
  cesarpintoi 437
  collinsae 437
  jequi 435–439
  jumbo 437
  prata 435–440
  robustus 435–440
  spilosoma 437
  spilurus 436–437
Paspalum fasciculatum 304

Passalus cornutus 166
Paulasquama callis 462
Pectenocypris korthausae 17, 19
Pectinantus parini 772
Pelamis 346
  platurus 346
Pelecus 14–15
  cultratus 17, 19
Pelias nikolskii 104
Pelobates 699
  cultripes 699
  fuscus 699
  syriacus 699–706
Pelobatidae 553
Pelodiscus sinensis 307
Pelophylax 103–110
  esculentus 104, 706
  lessonea 104, 706
  ridibundus 104, 106, 181, 706
Pentaclethra macroloba 224
Perca fluviatilis 107
Percidae 154–159, 354
Percina 625
  (Alvordius) 622
  copelandi 353–354
  crassa 622
  nevisense 622
  peltata 622–626
  phoxocephala 353
  rex 656
  roanoka 622, 656
  shumardi 353
  vigil 354–355
Percomorpha 23–30, 415–434
Percottus glenni 103–110
Periopthalmus 348
†*Perleidus* 587–589, 600
Peromyscus 806
Pterodroma cahow 191
Phaeognathus 266
  hubrichti 266
Pharyngostomum cordatum 105–107
Phelsuma 781
Phenacogaster 35
Phenacogrammus interruptus 16, 19, 315
Pholidophorus 587–589, 600
  bechei 586
Phosichthyidae 772
Photocorynus spiniceps 669
Phragmites 274
  australis 651, 700
Phrynohyas 634
  coracterias 639
  ingens 639
  latifasciata 639
  venulosa 706
Phrynosoma blainvillii 786
Phyllomedusa hypochondrialis 706
Phyllorhynchus 786
Physalaemus
  albonatus 706
  barrioi 798
  biligonigerus 706
  gracilis 706
Physalemus pustulosus 335
Physiculus parini 772
Physidae 477–484
Pimephales
  notatus 354
  promelas 164, 528
  vigilax 354
Pinus
  ponderosa 143
  sabineana 143
Pituophis 786
Plagiognathops 15
Planaltina 604–605, 610
Planorbidae 477–484
Planonasus parini 771–772
Plantago lanceolata 65
Platyberyx 415–434
  andriashevi 416, 418–420, 422–425, 432
  macropus 429
  mauli 415–416, 418, 423–424, 432
  opalescens 415–424, 432
  paucus 417–423, 432
  pietschi 417–424, 432
  rhyton 417–424, 432
  sp. 424
Platyceps 179
Platytroctidae 772
Plectranthias parini 772
Pleidae 105
Plethodon 504, 515, 676
  cinereus 179, 515, 676–683
  clemsonae 364
  metcalfi 364
Plethodontidae 515
Pleuroceridae 478–480
Pleurogenes claviger 106
Pleurogenoides medians 104, 106
†*Pliodetes* 582, 587–590, 600
Pneumonoeces
  asper 106
  variegatus 106
Podocnemididae 303–311
Podocnemis 111–120
  expansa 111–120, 307–308
  unifilis 111–120, 303–311
  vogli 111–120
Podostemaceae 460
Poecilia
  latipinna 80–96
  reticulata 204, 782
Poeciliidae 72–79
Polyipnus parini 772
Pomacentrus taeniometopon 533
Potamogeton 104
  sp. 692
Pristimantis educator 760
Pristobrycon striolatus 315
Procyon lotor 150
†*Propterus* 587–589, 600
Prosotocus confusus 104, 106
Prunus serotina 255–256
Psammophiidae 194–200
Psammophis 194
  schokari 198

Psammophylax 195
rhombeatus rhombeatus 194–200
Psellogrammus kennedyi 35
Pseudacris 673
crucifer 193
maculata 670–675
triseriata 671
Pseudechis porphyriacus 69
Pseudemys alabamensis 571
Pseudis paradoxa 706
Pseudobrama 15, 19
Pseudocurimata 630
Pseudohemiculter 19
Pseudolaubuca 15, 19
Pseudorasbora parva 17, 19
Pseudotriakidae 772
Pseudotsuga menziesii 143
Pseudoxygaster 15
Psilorhynchidae 18
Psilorhynchus
homaloptera 18–19
sucatio 18–19
Pteraclidae 425
Pteraclis 425
fasciatus 415, 427, 429
macropus 415, 424–425, 430
Petronia onobromoides 194
Pteronotropis hypselopterus 17, 19
Puntius
binotatus 410
ticto 16, 19
Pygidianops 441–444, 451, 641–642, 645
amphioxus 443, 447–449, 452, 645
cuao 452
eigenmanni 441, 452
magoi 443, 445, 452
Pygocentrus nattereri 315
Pyramodon parini 772
Pyrrhulina obermulleri 364
Python 179
Pythonidae 179

Quercus 255
garryana 143
kelloggii 143
rubra 677

Raiamas 14
bola 17, 19
senegalensis 17, 19
Rana
aurora 785
brevipoda 706
catesbeiana 278
dalmatina 702
draytonii 785
dybowskii 364
japonica 706
limnocharis 706
nigromaculata 706
pipiens 335
sylvatica 670, 717–722
tagoi 706
temporaria 703, 706
zografi 364
Ranidae 706
Rapateaceae 685
Rasbora cephalotaenia 17, 19
Rasborichthys 14–15
Rasborinae 14
Rasborinus 14–15, 19
Rasboroides vaterifloris 17, 19
Regalecidae 356
Regalecus 356
glesne 356
russellii 356
Retroculus xinguensis 449
Rhabdias
bufonis 106
fuscovenosus 105, 107
savagei 760
Rhacophorus
duboisi 220
hoanglienensis 220
Rhadinaea decorate 139
Rhamnus pallasii 65
Rhamphocetichthys savagei 760
Rhamphophryne 11
Rhinatrema
bivittatum 226
nigrum 224, 228
shiv 229
Rhinatrematidae 229
Rhinechis scalaris 139–140
Rhinella 11
marina 539–542, 548
Rhizophora mangle 81
Rhoadsia altipinna 35
Rhodeus ocellatus 16, 19
Rhondonanthus capillaceus 460
Rhoptropus afer 781
Rhynchorhamphus 770
Roeboides 35
Rutilus rutilus 107

Salamandra salamandra 514
Salmo
salar 353
trutta 353
Salmostoma 14–15
bacaila 17, 19
Salvadora 786
hexalepis 786
Sander canadensis 353
†*Sangiorgioichthys*
aldae 587–589, 599–600
sui 587–589, 599–600
Sarcocheilichthys variegatus 17, 19
Sarcoglanidinae 444, 645
Scaphiodontophis 132, 139
annulatus 139
Scaphiopus couchii 335
Scarabaeidae 105
Scardinius erythrophthalmus 353
Sceloporus
graciosus 785
grammicus disparilis 715
vandenbergianus 785
†*Scheenstia*
laevis 587–589, 600

mantelli 587–589, 600
maximus 587–589, 600
zappi 587–589, 600
Schilbeodes insignus 364
Schizodon 244
Schoenoplectus etuberculatus 657
Sciaenidae 740
Scinax acuminatus 706
Scincella schmidti 364
Scincidae 132, 800
Scomber scombrus 362
Scombropidae 24
Scutiger 213
(Aelurolalax) 213
chintingensis 213–215
Scyliorhinus 379
Scytopis hebes 639
Securicula 14–15
gora 17, 19
Seminatrix pygea 168
†*Semiolepis* 587–589, 600
brembanus 599
†Semionotidae 582–603
†Semionotiformes 582–603
†*Semionotus* 582–603
bergeri 587–590, 600
brembanus 599
capensis 586–590, 600
elegans 586–590, 600
fultus 599
kanabensis 582–603
micropterus 599
sp. 591, 599
tenuiceps 599
Semotilus atromaculatus 17, 19
Serpentes 132–141, 194–200
Serranidae 772, 798
Serrasalmus sp. 315
Sewellia lineolata 16, 19
†*Siemensichthys* 587–589, 600
macrocephalus 586
Siluriformes 16, 277, 396–402, 435–440, 577–581, 641–646
Sinibrama 19
Siphonopidae 229
Siphonops annulatus 229
Sistrurus 707
catenatus 507
c. catenatus 69, 97–102
Skiotocharax meizon 462
Skrjabinoeces similis 106
Sladenia 663
gardineri 663
shaefersi 660–665
Smiliogaster 13–14
Smiliogastri 13
Smiliogastrinae 13
Sparganium americanum 657
Spartina 736
patens 274
Spea hammondii 786
Sphaerodactylus savagei 760
Spintherobolus 35
Spiruridae 106
Squaliobarbidae 13–22
Squaliobarbus 13–22
curriculus 15, 18–19
Squamata 800
Stariophysi 800
Steatogenys elegans 232
Stegophilinae 645
Stegostenopos cryptogenes 232
Steindachnerina 630
Sterna antillarum browni 730
Sternoptychidae 772
Sternotherus carinatus 570
Stevardiinae 604
Stomiidae 772, 798
Storeria dekayi 167
Strigea
falconis 107
sphaerula 105–107
strigis 105–107
Strix occidentalis 324
Strongyloides
mirzai 107
spiralis 106
Sudis 41
gigas 40–41
pirarucu 40–41, 48
Symphysanodontidae 772
Symphysanodon parini 772
Synagrops 23–30
bellus 27
japonicus 25, 27

Tabanidae 105
Taeniopygia guttata 681
Tamiobatis 385
Tanakia limbata 16, 19
Tanichthys
albonubes 18–19
micagemmae 18–19
Tantilla hobartsmithi 786
Taricha
granulosa 335, 786
torosa 786
Tatia 396–402
carolae 396–399, 401
marthae 398–401
musaica 396, 400–401
Tekla 363
Telorchis assula 105, 107
Teleostei 13–22, 72–79, 356, 415–434, 441–453, 798
Terrapene
carolina carolina 570
ornata 570
Terrarana 179–180
Testudines 303–311
†*Tetragonolepis semicinctus* 600
Tetragonopterus
chalceus 35
ipanquianus 604–605, 610
Thalassoma bifasciatum 493–498
Thamnophis 132
butleri 139
cyrtopsis 786
elegans 198
marcianus 786
sauritus 132, 139

sirtalis 132, 139, 198
s. sirtalis 139
Thayeria sp. 616
Theloderma lateriticum 220
Thomisidae 105
Thunnus
alalunga 347
allisoni 363
Tilapia mariae 91
Tincinae 18
Tinca tinca 18–19
†*Tlayuamichin* 587–591, 600
Tometes lebaili 315
Tomistoma 517–526
schlegelii 518–520, 523–524
Toxabramis 15, 19
Trachemys scripta 61
Trachycephalus 634–640
atlas 634–635, 637
coriaceus 637
dibernardoi 634, 637
hadroceps 634–640
helioi 634–638
imitatrix 637
jordani 634–635, 637
lepidus 634, 638
mambaiensis 634–635, 638
mesophaeus 634, 638
nigromaculatus 634–635, 638
resinifictrix 634, 637–638
typhonius 634, 637–638
Trichiuridae 768
Trichogenes 443
Trichogeninae 444, 451
Trichomycteridae 441–453, 641–646
Trichomycterinae 444, 645
Trichomycterus
anhanga 645
hasemani 645
johnsoni 645
Trichopodus trichopterus 410
Trichoptera 478–479
Tricuspidalestes caeruleus 619
Tridentinae 645
Tridentopsis pearsoni 364
Trifolium pratense 65
Trimerisurus flavoviridis 167
Trimorphodon
lamba 786
lambda 786
Triticum aestivum 65
Truebella 11
Tylodelphys excavata 105–106
Typha
latifolia 103
sp. 700
Typhlobelus 441–453, 641–642, 645
auriculatus 441–453
cf. *macromycterus* 453
cuao 450
lundbergi 442–446, 450–453
guacamaya 442–446, 452
macromycterus 442–443, 449–453
sp. 450, 453
ternetzi 441–445, 449–450, 452–453
Typhlonectes compressicauda 229
Typhlonectidae 229
Tyttobrycon 35

Unidentata 206–212
Unionidae 478–480
Upeneus phillipsi 191
Urobatis jamaicensis 797
Urolophus cruciatus 787
Uromastyx hardwickii 179
Urosaurus nigricaudus 785
Urotheca 139
Uta stansburiana 798
Utricularia 81

Vaillantella maassi 16, 19
Vaillantellidae 16
Vandelliinae 444, 645
Vastres
agassizii 40
cuvieri 41, 48
mapae 41
pirarucu 48
Vertebrata 105
Vespidae 105
Vipera aspis 351
Viperidae 64, 707
Volodichthys parini 772

†*Watsonulus* 587–589, 600
eugnathoides 586

Xantusia
wigginsi 785
vigilis 785
Xenocypridina 19
Xenocypridinae 13
Xenocypridini 19
Xenocyprinidae 13–22
Xenocyprioides 13–22
parvulus 18–19
Xenocypris 13–22
argentea 15, 18–19
Xenopus laevis 351
Xeranthemum squarrosum 65
Xiphophorus 797
Xysticus nicholsi 805

Yaoshanicus 13–22

Zacco 13–22
Zoarcidae 772
Zootaca vipera 103
Zygophyllum morgsana 194

SUBJECT INDEX

Copeia 2013, Nos. 1–4

ACTIVITY, *Montivipera raddei* (spatial ecology; movement) 64–71; *Dendrophidion dendrophis, Mastigodryas bifossatus* (pseudoautotomy) 132–141; *Actinemys marmorata* (seasonal migration) 142–153; *Psammophylax r. rhombeatus* (spatial ecology; movement; sexual differences) 194–200; *Ambystoma* (*laterale-jeffersonianum* complex) (spatial ecology; habitat selection) 254–261; *Crotalus o. oreganus* (spatial ecology; movement; reproduction) 485–492; *Thalassoma bifasciatum* (larval settlement; spawning) 493–498; *Heterodon platirhinos* (road crossing; avoidance) 507–511; *Fundulus heteroclitus* (spawning in lab environment) 527–538; *Lithobates areolatus* (call surveying) 552–561; *Lithobates* (*catesbeianus, clamitans*) (sublethal effects of pesticides; predator–prey interaction) 691–698.

AGE, *Desmognathus* (*monticola, ocoee, quadramaculatus*) (age; reproduction; body size) 262–267; *Moxostoma* (*duquesnei, erythrurum*) (otolith) 390–395; *Thalassoma bifasciatum* (planktonic larval duration; otolith analysis) 493–498; *Etheostoma okaloosae* (longetivity; tagging method) 653–659.

AVOIDANCE, *Desmognathus* (*folkertsi, quadramaculatus*) (trophic interaction) 499–506; *Heterodon platirhinos* (road crossing) 507–511; *Lithobates* (*catesbeianus, clamitans*) (predators) 691–698.

BEHAVIOR, *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. (trophic interactions) 103–110; *Podocnemis* (*expansa, unifilis, vogli*) (nesting) 111–120; *Actinemys marmorata* (seasonal migration; flow regime) 142–153; *Etheostoma luteovinctum* (visual signals; male mating preference) 154–159; *Lampropholis delicata* (early hatching; predator escapement; sensory) 160–165; *Liolaemus chiliensis* (distress calls) 206–212; *Ambystoma* (*laterale-jeffersonianum* complex) (habitat selection) 254–261; *Micropterus salmoides* (predation; "dangerous prey" hypothesis) 277–283; *Ambystoma* (*talpoideum, maculatum*) (predation; cannibalism) 284–291; *Bufo melanostictus* (reproduction; plasma sex steroids) 332–337; *Graptemys geographica* (diet, sexual differences, environmental factors) 477–484; *Desmognathus* (*folkertsi, quadramaculatus*) (trophic interaction, competition) 499–506; *Desmognathus ocoee* (jumping; kinematics; scaling) 512–516; Crocodylia (long-distance signaling; evolutionary history) 517–526; *Rhinella marina* (stress response) 539–542; *Lithobates areolatus* (breeding calls) 552–561; *Hyphessobrycon* (*kayabi* n. sp.) (feeding) 616; *Trachycephalus* (*helioi* n. sp.) (probable breeding sites) 634–640; *Etheostoma okaloosae* (movement) 653–659; *Sladenia shaefersi* (*in-situ* observations; photos.) 660–665; *Centrophryne spinulosa* (reproduction; sexual parasitism) 666–669; *Plethodon cinereus* (assortative mating) 676–683; *Bothrops atrox* (habitat selection) 684–690.

BIODIVERSITY, *Incilius* (*majordomus* n. sp.) 8–12; *Hemigrammocharax* (*rubensteini* (n. sp.) (Congo basin) 31–37; *Arapaima agassizii* (validation) 38–51; *Characidium* (*papachibe* n. sp.) 52–57; *Gambusia* (*clarkhubbsi, krumholzi, speciosa*) (conservation status; species status; hybridization) 72–79; *Microcaecilia* (spp.) (Guianan Shield) 223–231; *Hypopygus* (species key) 236–237; *Hoplias microlepis* (morphologic/genetic divergence; haplotypes; environmental factors) 312–323; *Aspidoscelis* (*arizonae, pai, inornata llanuras*) (species diagnosis; validity) 366–377; *Tatia* (*carolae* n. sp., *marthae* n. sp.), (*musaica*; status re-evaluation) 396–402; *Oryzias* (Sulawesi; 19 valid spp.) 403–414; *Platyberyx, Caristius* (generic revision; 5 n. sp.) 415–434; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) 441–453; *Arapaima* (*leptosoma* n. sp., 5 nominal spp.) 470–476; *Doumea* (W African spp.) 580; †*Lophionotus* (*chinleana* n. sp.), †*L. kanabensis* n. comb. (redescription) 582–603; *Acrobrycon* (n. spp.; *A. tarijae* in synonymy) 604–611; *Cyphocharax* (highest in Curimatidae) 627–633; *Trachycephalus* (*helioi* n. sp.) 634–640; *Listrura* (*depinnai* n. sp.) 641–646.

BIOGEOGRAPHY, *Menidia audens* (invasive species; chronology; Tennessee/Cumberland R. systems) 292–302; *Hypomasticus* (*lineomaculatus* n. sp.) (river capture) 464–469; *Listrura* (*depinnai* n. sp.) (southernmost record, Glanapteryginae) 641–646; *Leuresthes tenuis* (northern habitat expansion; colonization events) 729–739.

BLOOD, *Sistrurus c. catenatus* (hematology; plasma biochemistry; heavy metals; disease) 97–102; *Bufo melanostictus* (plasma sex steroids; reproductive behavior) 332–337; *Ambystoma maculatum* (hematology; breeding M/F differences) 338–345.

CALLS [see also COMMUNICATION], *Liolaemus chiliensis* (distress; sexual variation; spectrograms/oscillograms) 206–212; Crocodylia (signal components; long-distance signaling; evolutionary history) 517–526; *Lithobates areolatus* (breeding survey method) 552–561; *Trachycephalus* (*helioi* n. sp.) (described; single vocal sac) 634–640.

CANNIBALISM, *Ambystoma* (*talpoideum, maculatum*) (conspecific/congeneric) 284–291.

CHEMICAL CUES, *Ambystoma* (*laterale-jeffersonianum* complex) (habitat selection) 254–261; *Desmognathus* (*folkertsi, quadramaculatus*) (trophic interaction, avoidance) 499–506; *Plethodon cinereus* (assortative mating; male scent, fecal odors) 676–683.

CHONDROCRANIUM, *Lamna* (*ditropis, nasus*) (description; photos.) 378–389.

COLONIZATION, *Leuresthes tenuis* (range extension; disjunct events) 729–739.

COLORATION [see also PIGMENTATION], *Incilius* (*majordomus* n. sp.) (sexual dichromatism; in life/preservation) 8–12; *Amioides* (*polyacanthus* n. comb.) (perimortem/in alcohol; figs.) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) (in life/alcohol; figs.) 31–37; *Arapaima agassizii* (based on original painting) 38–51; *Characidium* (*papachibe* n. sp.) (in ethanol; diagnostic pattern) 52–57; *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) (dewlap; species variation; pterins, carotenoids) 201–205; *Oreolalax* (*sterlingae* n. sp.) (preservative/alcohol; figs.) 213–222; *Microcaecilia* (*savagei* n. sp.), *Epicrionops niger* (in life; figs.) 223–231; *Hypopygus* (*benoneae* n. sp.) (in alcohol; fig.) 232–237; *Leporinus* (*moralesi, parvulus* n. sp.) (in life; figs.) 238–247; *Aspidoscelis* (*arizonae, pai, inornata llanuras*) (diagnostic patterns; figs.) 366–377; *Tatia* (*carolae* n. sp., *marthae* n. sp.) (in alcohol; figs.) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (in life/alcohol; color photos.; distinguishing character) 403–414; *Parotocinclus* (*jequi* n. sp.) (in alcohol; fig.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) (in life/alcohol; figs.) 441–453; *Characidium* (*amaila* n. sp.) (diagnostic; in life/alcohol; figs.) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) (in alcohol; figs.) 464–469; *Arapaima* (*leptosoma* n. sp.) (in alcohol) 470–476; *Doumea* (*skeltoni* n. sp.) (in alcohol; fig.) 577–581; *Acrobrycon* (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) (in alcohol; figs.) 604–611; *Hyphessobrycon* (reticulate-pattern clade) (*kayabi* n. sp.) (in life/alcohol; fig.) 612–621; *Percina peltata* (population variation) 622–626; *Cyphocharax* (*aninha* n. sp.) (in ethanol; pigmentation character; fig.) 627–633; *Trachycephalus* (*helioi* n. sp.) (in life; holotype; figs.) 634–640; *Listrura* (*depinnai* n. sp.) (in life/alcohol; figs.) 641–646.

COMMUNICATION [see also CALLS], *Liolaemus chiliensis* (distress calls) 206–212; Crocodylia (long-distance signaling) 517–526.

COMMUNITY, *Cichlasoma urophthalmus* (invasive; effect on native fish) 80–96; *Perccottus glenii, Natrix* (*natrix, tessellata*), (*Pelophylax* spp.) (trophic interactions) 103–110; *Lithobates clamitans* (environmental factors) 248–253; *Ambystoma* (*talpoideum, maculatum*) (intraguild predation; cannibalism) 284–291; *Lithobates* (*catesbeianus, clamitans*) (predator–prey interactions; sublethal pesticides) 691–698.

© 2013 by the American Society of Ichthyologists and Herpetologists 10.1643/OT-13-138

COMPETITION, *Menidia audens, Labidesthes sicculus* (invasive species; population decline) 292–302.

COMPETITION–PREDATION HYPOTHESIS, *Desmognathus* (*folkertsi, quadramaculatus*) 499–506.

CONSERVATION, *Arapaima agassizii* (status unknown) 38–51; *Macrochelys temminckii* (harvesting; population; recovery) 58–63; *Montivipera raddei* (spatial ecology; human-modified landscape) 64–71; *Gambusia krumholzi* (populations; management units) 72–79; *Podocnemis* (*expansa, unifilis, vogli*) (suggested methods) 111–120; *Glyptemys muhlenbergii* (population gene flow; regional approach) 324–331; *Aspidoscelis arizonae* (species status) 366–377; *Arapaima* (*leptosoma* n. sp.) (restricted distribution) 470–476; anuran larvae (7 spp.) (salinity tolerances, survival) 543–551; *Lithobates areolatus* (threatened species) 552–561; *Lithobates belandieri, Incilius nebulifer, Acris crepitans* (effects of salinity increases) 562–566; *Percina peltata* (isolated population; morphological distinctness) 622–626; *Chrosomus* sp. cf. *saylori* (fragmented populations) 750–756.

"DANGEROUS PREY" HYPOTHESIS, *Micropterus salmoides* (feeding; spination avoidance) 277–283.

DEEP-SEA, *Amioides* (*polyacanthus* n. comb.) (Indo-Pacific; 77–276 m) 23–30; *Platyberyx* (*andriashevi, mauli, opalescens, paucus* n. sp., *pietschi* n. sp., *rhyton* n. sp.), *Caristius* (*digitus* n. sp., *fasciatus, macropus, meridionalis* n. sp.) (descriptions; distributions; maps; generic revision) 415–434; *Sladenia shaefersi* (*in-situ* observations; 689–1200 m depth) 660–665; *Centrophryne spinulosa* (sexual parasitism) 666–669.

DEFENSE MECHANISMS, *Desmognathus ocoee* (jumping behavior) 512–516; *Crotalus lepidus morulus* (venom, composition & activity) 707–716.

DENTITION, *Arapaima agassizii* (basibranchial toothplate; diagnostic; fig.) 38–51; *Hyphessobrycon* (*kayabi* n. sp.) 612–621.

DEVELOPMENT, *Moxostoma duquesnei* (ontogeny; pigmentation; morphology) 121–126; *Lithobates clamitans* (environmental factors) 248–253; *Podocnemis unifilis* (embryonic development; abiotic environmental factors) 303–311; *Rana sylvatica* (environmental factors; growth, locomotor performance) 717–722; *Aplodinotus grunniens* (morphologic variation) 740–749.

DIET, *Graptemys geographica* (prey choice; sexual differences; environmental factors) 477–484; *Hyphessobrycon* (*kayabi* n. sp.) (stomach contents) 612–621; *Plethodon cinereus* (fecal odor cue) 676–683; *Chrosomus* sp. cf. *saylori* (atypical trophic niche) 750–756.

DISEASE, *Sistrurus c. catenatus* (*Chrysosporium* fungus; heavy metals; health survey) 97–102.

DISTRIBUTION, *Incilius* (*majordomus* n. sp.) (Pacific slope, Panama; map) 8–12; *Amioides* (*polyacanthus* n. comb.) (Indo-Pacific) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) (Lekoli R., Congo; map) 31–37; *Arapaima agassizii* (holotype only: lowlands, Brazilian Amazon) 38–51; *Characidium* (*papachibe* n. sp.) (type loc. only: tributary, Rio Aruã, Brazil; map) 52–57; *Gambusia krumholzi* (restricted populations; Texas, Mexico; map) 72–79; *Cichlasoma urophthalmus* (invasive species; estuarine Everglades; map) 80–96; *Oreolalax* (range extension), (*O. sterlingae* n. sp.) (Vietnam; map) 213–222; *Microcaecilia* (*savagei* n. sp.) (type loc. only; Iwokrama Forest), *Epicrionops niger* (range extension), (Guyana; map) 223–231; *Hypopygus* (*benoneae* n. sp.) (Rio Anapu, Brazil; map) 232–237; *Leporinus* (*moralesi, parvulus* n. sp.) (Amazon basin, Brazil; map) 238–247; *Ambystoma* (*laterale-jeffersonianum* complex) (habitat preference/avoidance) 254–261; *Menidia audens* (invasive species; Tennessee/Cumberland R. systems; maps; chronology) 292–302; *Hoplias microlepis* (Guayas R. drainage; map) 312–323; *Glyptemys muhlenbergii* (range map; conservation) 324–331; *Aspidoscelis arizonae* (SE Arizona; map) 366–377; *Tatia* (*carolae* n. sp.) (type loc. only, Cuyuni R.), (*marthae* n. sp.) (type loc. only, Rio Caura) 390–395; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (SE Sulawesi; map) 403–414; *Platyberyx* (6 spp.), *Caristius* (4 spp.) (worldwide; maps) 415–434; *Parotocinclus* (*jequi* n. sp.) (type loc. only, Rio Jequitinhohoa; map) 435–440; *Typhlobelus* (*auriculatus* n. sp.) (type loc. only, lower Rio Xingu; map) 441–453; *Characidium* (*amaila* n. sp.) (upper Kuribrong R., Guyana; map) 454–463; *Hypomasticus* (*despaxi, lineomaculatus* n. sp.) (Guyana-Amazon provinces; map) 464–469; *Arapaima* (*leptosoma* n. sp.) (Solimoes R., Brazil) 470–476; *Doumea* (*skeltoni* n. sp.) (NW Kouilou R. system; map) 577–581; †*Lophionotus* (*chinleana* n. sp., *kanabensis* n. comb.) (S-SE Utah fossil sites; map) 582–603; *Acrobrycon* (W Rio Amazonas thru NW Rio de la Plata; map) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) (type loc., Rio Teles Pires; map) 612–621; *Percina peltata* (isolated population; map) 622–626; *Cyphocharax* (*aninha* n. sp.) (Rio Mopeco; map) 627–633; *Trachycephalus* (*helioi* n. sp.) (restricted, Amazon R.; comparison to *T. hadroceps*; map) 634–640; *Listrura* (coastal Atlantic Forest, Brazil; map), (*depinnai* n. sp.) (type loc. only, Laguna des Patos) 641–646; *Sladenia shaefersi* (No. Atlantic, Gulf of Mexico; map) 660–665; *Pelobates syriacus* (Danube Delta Biosphere Reserve) 699–706; *Leuresthes tenuis* (northern range expansion, disjunct events; map) 729–739.

ECOLOGY, *Incilius* (*majordomus* n. sp.) (distribution; habitat; sexual dichromatism) 8–12; *Amioides* (*polyacanthus* n. comb.) (distribution; habitat) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) (distribution; habitat) 31–37; *Characidium* (*papachibe* n. sp.) (distribution; habitat; sexual dimorphism) 52–57; *Macrochelys temminckii* (population recovery; sex ratio; survivorship; adult: juvenile ratio) 58–63; *Montivipera raddei* (spatial ecology; movement; habitat) 64–71; *Cichlasoma urophthalmus* (invasive species; prey) 80–96; *Sistrurus c. catenatus* (fungal disease; heavy metal; health survey) 97–102; *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. (trophic interactions; prey; helminth parasites) 103–110; *Podocnemis* (*expansa, unifilis, vogli*) (harvesting; sex ratio; size) 111–120; *Dendrophidion dendrophis, Mastigodryas bifossatus* (pseudoautotomy) 132–141; *Actinemys marmorata* (seasonal migration; flow regime; life history strategies) 142–153; *Etheostoma luteovinctum* (mating preference; visual cues) 154–159; *Lampropholis delicata* (early hatching; predation escapement; sensory; survival) 160–165; *Psammophylax r. rhombeatus* (spatial ecology; movement; M/F home ranges) 194–200; *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) (dewlap coloration) 201–205; *Liolaemus chiliensis* (distress calls; sexual variation) 206–212; *Oreolalax* (*sterlingae* n. sp.) (distribution; habitat; sexual dimorphism) 213–222; *Microcaecilia* (*savagei* n. sp.) (distribution; habitat), *Epicrionops niger* (distribution) 223–231; *Hypopygus* (*benoneae* n. sp.) (distribution; habitat) 232–237; *Leporinus* (*moralesi, parvulus* n. sp.) (distribution; habitat) 238–247; *Lithobates clamitans* (tadpole growth/survivorship; environmental factors) 248–253; *Ambystoma* (*laterale-jeffersonianum* complex) (habitat selection; spatial distribution; environmental factors) 254–261; *Desmognathus* (*monticola, ocoee, quadramaculatus*) (age, reproduction; fecundity; size; mortality rate) 262–267; *Alligator mississippiensis* (reproductive allometry; egg/clutch/nest size; habitat quality) 268–276; *Micropterus salmoides* (feeding; "dangerous prey" hypothesis) 277–283; *Ambystoma* (*talpoideum, maculatum*) (intraguild predation; cannibalism; paedomorphs, overwintering larvae) 284–291; *Menidia audens, Labidesthes sicculus* (invasive species; competition; population decline) 292–302; *Podocnemis unifilis* (embryonic development; abiotic environmental factors; nesting sites; nest characteristics; hatching success; sex ratio; survival; reproduction) 303–311; *Hoplias microlepis* (morphologic/genetic divergence; habitat alteration; environmental factors; haplotypes) 312–323; *Glyptemys muhlenbergii* (gene flow, fragmented populations; conservation) 324–331; *Bufo melanostictus* (reproductive behavior; plasma sex steroids) 332–337; *Ambystoma maculatum* (sexual dimorphism; hematology; visceral mass; reproduction) 338–345; *Aspidoscelis arizonae* (distribution; habitat) 366–377; *Moxostoma* (*duquesnei, erythrurum*) (age; growth) 390–395; *Tatia* (*carolae* n. sp., *marthae* n. sp.) (distribution) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (distribution; habitat; reproduction) 403–414; *Platyberyx, Caristius* (distribution) 415–434; *Parotocinclus* (*jequi* n. sp.) (distribution; habitat; sexual dimorphism) 435–440; *Typhlobelus* (*auriculatus* n. sp.) (distribution; habitat; sexual dimorphism; endangered) 441–453; *Characidium* (*amaila* n. sp.) (distribution; habitat; rheophilic) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) (distribution) 464–469; *Arapaima* (*leptosoma* n. sp.) (distribution) 470–476; *Graptemys geographica* (diet; sexual differences) 477–484; *Crotalus o. oreganus* (spatial ecology; home range; reproduction) 485–492;

Thalassoma bifasciatum (planktonic larval duration; settlement; spawning) 493–498; *Desmognathus* (*folkertsi, quadramaculatus*) (trophic interaction, competition) 499–506; *Heterodon platirhinos* (movement) 507–511; *Desmognathus ocoee* (jumping behavior; defense mechanism) 512–516; Crocodylia (long-distance signaling) 517–526; *Fundulus heteroclitus* (spawning in lab environment; egg production; gonad size) 527–538; *Rhinella marina* (stress response; corticosterone levels) 539–542; anuran larvae (7 spp.) (salinity tolerance; survival) 543–551; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (lethal/sublethal effects of salinity) 562–566; *Doumea* (*skeltoni* n. sp.) (distribution; habitat) 577–581; *Acrobrycon* (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) (distribution; habitat; 2nd sexual dimorphism) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) (distribution; habitat; sexual dimorphism; diet; feeding) 612–621; *Percina peltata* (distinct population) 622–626; *Cyphocharax* (*aninha* n. sp.) (distribution; habitat) 627–633; *Trachycephalus* (*helioi* n. sp.) (distribution; habitat; calls; breeding site) 634–640; *Listrura* (*depinnai* n. sp.) (distribution; habitat) 641–646; *Anaxyrus* (=*Bufo*) *fowleri* (sex ratio; breeding sites; population/abundance) 647–652; *Etheostoma okaloosae* (movement; longetivity; population dynamics) 653–659; *Sladenia shaefersi* (distribution; habitat; mating behavior) 660–665; *Centrophryne spinulosa* (sexual parasitism) 666–669; *Pseudacris maculata* (marking methods, survival) 670–675; *Plethodon cinereus* (assortative mating; scent/diet cues) 676–683; *Bothrops atrox* (habitat selection; movement) 684–690; *Lithobates* (*catesbeianus, clamitans*) (predator–prey interactions; sublethal effects of pesticides) 691–698; *Pelobates syriacus* (reproductive strategies) 699–706; *Crotalus lepidus morulus* (infraspecific venom variation) 707–716; *Bothrops atrox* (environmental factors & development) 717–722; *Graptemys caglei* (genetic variation between populations; gene flow) 723–728; *Leuresthes tenuis* (habitat expansion; no./so. population comparisons) 729–739; *Aplodinotus grunniens* (environmental factors & morphology) 740–749; *Chrosomus* sp. cf. *saylori* (diet preference; trophic niche) 750–756.

EGGS, *Moxostoma duquesnei* (size; hatching) 121–126; *Lampropholis delicata* (early hatching; predator escapement; sensory) 160–165; *Oreolalax* (*sterlingae* n. sp.) (coloration; size) 213–222; *Desmognathus* (*monticola, ocoee, quadramaculatus*) (follicle number; body-size) 262–267; *Alligator mississippiensis* (size; reproductive allometry) 268–276; *Podocnemis unifilis* (clutch size; hatching success) 303–311; *Fundulus heteroclitus* (production over time) 527–538; *Pelobates syriacus* (clutch size; egg size, number) 699–706; *Leuresthes tenuis* (No./So. populations; clutch volume; egg diameter) 729–739.

ENDANGERED SPECIES, *Arapaima agassizii* (restricted distribution) 38–51; *Gambusia krumholzi* (restricted distribution; management units) 72–79; *Podocnemis* (*expansa, unifilis, vogli*) (harvesting) 111–120; *Labidesthes sicculus* (population decline; invasive species; chronology) 292–302; *Glyptemys muhlenbergii* (fragmented populations; gene flow) 324–331; *Typhlobelus* (*auriculatus* n. sp.) (threat, dam construction) 441–453; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (increasing salinity levels) 562–566; *Etheostoma okaloosae* (imperiled) 653–659.

ENDEMIC/ENDEMISM, *Oryzias* (*asinua* n. sp.) (type loc. only) 403–414; *Typhlobelus* (*auriculatus* n. sp.) (type loc. only; endangered habitat) 441–453; *Characidium* (*amaila* n. sp.) (upper Kuribrong R., Guyana) 454–463; *Acrobrycon* (*starnesi* n. sp.) (type loc. only, Rio Thyumayu), (*ortii* n. sp.) (type loc. only, Rio Pilcomayo) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) (type loc. only, Rio Teles Pires) 612–621; *Cyphocharax* (*aninha* n. sp.) (type loc. only, Rio Mopeco) 627–633.

ENVIRONMENTAL FACTORS (Abiotic/Biotic), *Actinemys marmorata* (flow regime; behavioral/physical responses) 142–153; *Lithobates clamitans* (tadpole survivorship; invasive species/fertilizer) 248–253; *Ambystoma* (*laterale-jeffersonianum* complex) (habitat selection; chemical cues; leaf litter) 254–261; *Alligator mississippiensis* (habitat quality; reproductive allometry) 268–276; *Podocnemis unifilis* (nests; embryonic development; hatching success) 303–311; *Hoplias microlepis* (morphologic/genetic divergence; habitat alteration; impoundments) 312–323; *Moxostoma* (*duquesnei, erythrurum*) (otolith annulus formation) 390–395; *Graptemys geographica* (diet, sexual differences) 477–484; *Crotalus o. oreganus* (home range, movements, sexual size differences) 485–492; *Heterodon platirhinos* (road crossing) 507–511; *Fundulus heteroclitus* (spawning in lab environment; lunar cycle; latitudinal gradient) 527–538; *Rhinella marina* (stress response; toe-clipping vs. handling) 539–542; anuran larvae (7 spp.) (salinity level tolerance; survival) 543–551; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (lethal/sublethal salinity levels) 562–566; *Pseudacris maculata* (marking methods, survival) 670–675; *Bothrops atrox* (landscape characteristics) 684–690; *Lithobates* (*catesbeianus, clamitans*) (sublethal pesticides; predator–prey interactions) 691–698; *Rana sylvatica* (canopy cover & development) 717–722; *Aplodinotus grunniens* (morphological variation) 740–749.

EVOLUTION, *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) (pigmentation variation) 201–205; *Aspidoscelis arizonae* (morphological divergence) 366–377; Crocodylia (long-distance signaling) 517–526.

EXTINCTION, *Typhlobelus* (*auriculatus* n. sp.) (threat, habitat destruction) 441–453.

FECUNDITY, *Desmognathus* (*monticola, ocoee, quadramaculatus*) (body size; age; reproduction) 262–267; *Alligator mississippiensis* (clutch size; reproductive allometry) 268–276; *Pelobates syriacus* (clutch size) 699–706.

FEEDING, *Cichlasoma urophthalmus* (invasive species; prey) 80–96; *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. (trophic interactions; parasite transmission) 103–110; *Micropterus salmoides* (prey selection; "dangerous prey" hypothesis) 277–283; *Ambystoma* (*talpoideum, maculatum*) (intraguild predation; cannibalism) 284–291; *Graptemys geographica* (sexual differences; prey selection) 477–484; *Hyphessobrycon* (*kayabi* n. sp.) (behavior) 612–621.

FOSSIL, *Lamna* (fossil rostral cartilage) 378–389; †*Lophionotus* (redescription), (*chinleana* n. sp., *kanabensis* n. comb.) (descriptions; osteology; figs.; horizons) 582–603.

FUNGUS, *Sistrurus c. catenatus* (*Chrysosporium* prevalence) 103–110.

GENETICS, *Oxygaster*, Oxygastrinae (mitochondrial/nuclear DNA; placement; taxonomic status; maximum likelihood topology) 8–12; *Gambusia* (*krumholzi, clarkhubbsi, speciosa*) (conservation genetics; species status; hybridization) 72–79; *Oreolalax* (*sterlingae* n. sp.) (mtDNA) 213–222; *Hoplias microlepis* (mtDNA; haplotypes; population divergence) 312–323; *Glyptemys muhlenbergii* (microsatellite DNA; fragmented populations; gene flow) 324–331; *Oryzias* (*woworae* sp. gr.) (mtDNA; maximum parsimony & Bayesian inference trees) 403–414; *Graptemys caglei* (gene flow; microsatellite DNA) 723–728.

GEOGRAPHIC LOCALITY,
Alabama, *Menidia audens, Labidesthes sicculus* 292–302.
Arizona, *Aspidoscelis* (*pai, arizonae*) 366–377.
Arkansas, *Macrochelys temminckii* 58–63.
Argentina, *Acrobrycon ipanquianus* 604–611.
Armenia (Republic of), *Montivipera raddei* 64–71.
Atlantic Ocean, *Platyberyx* (*andriashevi, mauli, opalescens, paucus* n. sp., *pietschi* n. sp.), *Caristius fasciatus* 415–434; *Sladenia shaefersi* 660–665; *Centrophryne spinulosa* 666–669.
Australia, (NSW) *Lampropholis delicata* 160–165.
Bolivia, *Acrobrycon* (*starnesi* n. sp., *ortii* n. sp., *ipanquianus*) 604–611.
Brazil, *Arapaima agassizii* 38–51; *Characidium* (*papachibe* n. sp.) 52–57; *Dendrophidion dendrophis, Mastigodryas bifossatus* 132–141; *Hypopygus* (*benoneae* n. sp.) 232–237; *Leporinus* (*moralesi, parvulus* n. sp.) 238–247; *Podocnemis unifilis* 303–311; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) 441–453; *Hypomasticus* (*lineomaculatus* n. sp.) 464–469; *Arapaima* (*leptosoma* n. sp.) 470–476; *Hyphessobrycon* (*kayabi* n. sp.) 612–621; *Cyphocharax* (*aninha* n. sp.) 627–633; *Trachycephalus* (*helioi* n. sp.) 634–640; *Listrura* (*depinnai* n. sp.) 641–646; *Bothrops atrox* 684–690.
California, *Actinemys marmorata* 142–153; *Crotalus o. oreganus* 485–492; *Leuresthes tenuis* 729–739.
Canada, (Ont.) *Moxostoma duquesnei* 121–126; (Ont.) *Heterodon platirhinos* 507–511; (N.B.) *Fundulus heteroclitus* 527–538; (Ont.) *Anaxyrus* (=*Bufo*) *fowleri* 653–659.
Chile, *Liolaemus chiliensis* 206–212.
Congo (Republic of), *Hemigrammocharax* (*rubensteini* n. sp.) 31–37; *Doumea* (*skeltoni* n. sp.) 577–581.

Cuba, *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) 201–205.
Ecuador, *Hoplias microlepis* 312–323.
Florida, *Cichlasoma urophthalmus, Cyprinodon variegatus, Fundulus confluentus, Gambusia holbrooki, Lepomis* spp., *Menidia peninsulae, Microgobius gulosus, Poecilia latipinna* 80–96; *Hyla* (*cinerea, squirella*), *Lithobates* (*catesbeianus, sphenocephalus*), *Anaxyrus terrestris, Gastrophryne carolinensis, Osteopilus septentrionalis* 543–551.
Georgia, *Desmognathus* (*folkertsi, quadramaculatus*) 499–506.
Guyana, *Microcaecilia* (*savagei* n. sp.), *Epicrionops niger* 223–231; *Tatia* (*carolae* n. sp.) 396–402; *Characidium* (*amaila* n. sp.) 454–463.
Illinois, *Sistrurus c. catenatus* 97–102.
India, *Bufo melanostictus* 332–337.
Indian Ocean, *Platyberyx* (*andriashevi*), *Caristius* (*digitus* n. sp.) 415–434.
Indiana, *Ambystoma maculatum* 338–345; *Lithobates areolatus* 552–561; *Aplodinotus grunniens* 740–749.
Indo-Pacific, *Amioides* (*polyacanthus* n. comb.) 23–30.
Indonesia, (Sulawesi) *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) 403–414.
Kentucky, *Ambystoma* (*talpoideum, maculatum*) 284–291; *Menidia audens, Labidesthes sicculus* 292–302.
Louisiana, *Alligator mississippiensis* 268–276.
Maryland, *Graptemys geographica* 477–484; *Percina peltata* 622–626.
Massachusetts, *Glyptemys muhlenbergii* 324–331; *Fundulus heteroclitus* 527–538.
Mexico, *Gambusia* (*krumholzi, speciosa*) 72–79.
Michigan, *Ambystoma* (*laterale-jeffersonianum* complex) 254–261; *Rana sylvatica* 717–722.
Missouri, *Moxostoma* (*duquesnei, erythrurum*) 390–395.
New Mexico, *Aspidoscelis inornata llanuras* 366–377.
New York, *Glyptemys muhlenbergii* 324–331.
North Carolina, *Desmognathus* (*monticola, ocoee, quadramaculatus*) 262–267; *Menidia audens, Labidesthes sicculus* 292–302.
Ohio, *Lithobates clamitans, Gambusia affinis* 248–253; *Plethodon cinereus* 676–683.
Pacific Ocean, *Platyberyx* (*andriashevi, paucus* n. sp., *pietschi* n. sp., *rhyton* n. sp.), *Caristius* (*digitus* n. sp., *macropus*) 415–434.
Panama, *Incilius* (*majordomus* n. sp., *peripatetes*) 8–12.
Pennsylvania, *Percina peltata* 622–626.
Peru, *Acrobrycon ipanquianus* 604–611.
Roumania, *Pelobates syriacus* 699–706.
Russia, *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. 103–110.
South Africa, *Psammophylax r. rhombeatus* 194–200.
South Carolina, *Desmognathus* (*folkertsi, quadramaculatus*) 499–506.
Tennessee, *Menidia audens, Labidesthes sicculus* 292–302; *Tatia* (*marthae* n. sp.) 396–402.
Texas, *Gambusia* (cf. *clarkhubbsi, speciosa*) 72–79; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* 562–566; *Graptemys caglei* 723–728.
Utah, †*Lophionotus* (*chinleana* n. sp., *kanabensis* n. comb.) 582–603.
Venezuela, *Podocnemis* (*expansa, unifilis, vogli*) 111–120; *Tatia* (*marthae* n. sp.) 396–402.
Vietnam, *Oreolalax* (*sterlingae* n. sp.) 213–222.
Virgin Islands (U.S.), *Thalassoma bifasciatum* 493–498.
Virginia, *Menidia audens, Labidesthes sicculus* 292–302; *Fundulus heteroclitus* 527–538; *Chrosomus* sp. cf. *saylori* 750–756.

GROWTH, *Lithobates clamitans* (environmental factors) 248–253; *Moxostoma* (*duquesnei, erythrurum*) (otolith; longetivity) 390–395; *Pelobates syriacus* (reproductive strategy) 699–706; *Rana sylvatica* (effects of canopy cover) 717–722; *Aplodinotus grunniens* (allometry; environmental factors; variation) 740–749.

HABITAT, *Incilius* (*majordomus* n. sp.) (montane rainforest) 8–12; *Amioides* (*polyacanthus* n. comb.) (deepwater, 77–276 m; drop-offs/caves) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) (small tributaries) 31–37; *Characidium* (*papachibe* n. sp.) (shallow water, tributaries) 52–57; *Montivipera raddei* (human-modified; preferences; home range size) 64–71; *Cichlasoma urophthalmus* (invasive; estuarine Everglades) 80–96; *Perccottus glenii* (invasive; shallow lake) 103–110; *Psammophylax r. rhombeatus* (coastal shrub; home range) 194–200; *Oreolalax* (*sterlingae* n. sp.) (bamboo forest, elev. 2900 m; photo.) 213–222; *Microcaecilia* (*savagei* n. sp.) (leaf-litter; type loc. only) 223–231; *Hypopygus* (*benoneae* n. sp.) (rainforest stream) 232–237; *Leporinus* (*parvulus* n. sp.) (rocky pools/rapids) 238–247; *Ambystoma* (*laterale-jeffersonianum* complex) (leaf litter; preference/avoidance; chemical cues) 254–261; *Alligator mississippiensis* (quality effect on nest/clutch) 268–276; *Glyptemys muhlenbergii* (fragmented; conservation; gene flow) 324–331; *Oreolalax* (*sterlingae* n. sp.) (bamboo forest, elev. 2900 m; photo) 213–222; *Aspidoscelis arizonae* (SE Arizona; photos.) 366–377; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (fw, type loc.; photos.) 403–414; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) (psammophilic; habitat destruction; photo.) 441–453; *Characidium* (*amaila* n. sp.) (rapids; photos.) 454–463; *Graptemys geographica* (abiotic/biotic alterations & diet) 477–484; anuran larvae (7 spp.) (fw coastal wetlands; salinity level) 543–551; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (increasing salinity levels) 562–566; †*Lophionotus* (*chinleana* n. sp.) (fluvial-lacustrine; Chinle Fm.), †*L. kanabensis* n. comb. (lacustrine; Moenave Fm.) 582–603; *Hyphessobrycon* (*kayabi* n. sp.) (terre-firme lake) 612–622; *Trachycephalus* (*helioi* n. sp.) (canopy, tropical rain forest) 634–640; *Listrura* (*depinnai* n. sp.) (swamp, black water) 641–646; *Sladenia shaefersi* (hard bottom/coral rubble; 689–1200 m depth) 660–665; *Bothrops atrox* (riparian/non-riparian, selection) 684–690; *Rana sylvatica* (canopy growth & development) 717–722.

HARVESTING, *Macrochelys temminckii* (population recovery; survivorship) 58–63; *Podocnemis* (*expansa, unifilis, vogli*) (conservation efforts) 111–120.

HATCHLINGS, *Podocnemis unifilis* (hatching success; sex ratio; environmental factors) 303–311.

HELMINTHS, *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax esculentus* (trophic interactions; lists, prevalence/abundance/infection sites) 103–110.

HISTOLOGY, *Gasterosteus aculeatus* (cartilage development; staining technique) 127–131; *Dendrophidion dendrophis, Mastigodryas bifossatus* (caudal morphology) 132–141.

HISTORICAL PERSPECTIVES, H. G. Dowling, Jr. 166–172; *Copeia* (journal origin) 189–193, 361–365; J. M. Savage 757–767.

HISTORICAL RECORD, *Menidia audens* (invasive species; Tennessee/Cumberland R. systems; table) 292–302; *Aspidoscelis arizonae* (taxonomy) 366–377.

HOME RANGE, *Montivipera raddei* (M/F movements; minimum convex polygons; figs.) 64–71; *Psammophylax r. rhombeatus* (M/F movements; minimum convex polygons; figs.) 194–200; *Crotalus o. oreganus* (sexual differences; home range; method comparisons) 485–492.

HORMONES, *Bufo melanostictus* (plasma sex steroids) 332–337; *Rhinella marina* (corticosterone levels; stress response) 539–542.

HYBRIDIZATION/HYBRIDS, *Gambusia* (*krumholzi, speciosa*) (mitochondrial/microsatellite DNA; haplotypes) 72–79.

INVASIVE SPECIES, *Cichlasoma urophthalmus* (estuarine Everglades; trophic interactions) 80–96; *Perccottus glenii* (Russian lake; trophic interactions; parasite transmission) 103–110; *Gambusia affinis, Lithobates clamitans* (trophic interaction) 248–253; *Menidia audens* (Tennessee/Cumberland R. systems; chronology) 292–302; *Osteopilus septentrionalis* (salinity tolerance; invasive potential) 543–551.

LARVAE [see also TADPOLES], *Moxostoma duquesnei* (development; morphometrics; congeneric comparisons) 121–126; *Lithobates clamitans* (growth; survivorship; environmental factors) 248–253; *Thalassoma bifasciatum* (planktonic stage duration; settlement) 493–498; anurans (7 spp.) (salinity tolerance; survival) 543–551.

LATEROSENSORY PORES, *Typhlobelus* (*auriculatus* n. sp.) (absence; unique) 441–453.

LIFE HISTORY [see also NATURAL HISTORY], *Macrochelys temminckii* (harvesting; survivorship; sex ratio; adult:juvenile ratio) 58–63; *Actinemys marmorata* (flow regime/strategies; seasonal migration) 142–153; *Lampropholis delicata* (early hatching; predator escapement; survivorship) 160–165; *Desmognathus* (*monticola, ocoee, quadramaculatus*) (body size; fecundity; age; reproduction; mortality rate) 262–267; *Bufo melanostictus* (reproductive behavior; plasma sex steroids) 332–337; *Moxostoma* (*duquesnei, erythrurum*) (age; growth) 390–395; *Centrophryne*

spinulosa (sexual parasitism) 666–669; *Rana sylvatica* (development & canopy cover) 717–722; *Chrosomus* sp cf. *saylori* (ontogeny; growth) 750–756.

LOCOMOTION, *Rana sylvatica* (effects of canopy cover) 717–722.

MERISTICS/MORPHOMETRICS, *Incilius* (*majordomus* n. sp.) 8–12; *Hemigrammocharax* (*rubensteini* n. sp.) 31–37; *Arapaima agassizii* (descriptions; congeneric comparisons) 38–51; *Characidium* (*papachibe* n. sp.) 52–57; *Moxostoma duquesnei* (larval stages; congeneric comparisons) 121–126; *Actinemys marmorata* (body size) 142–153; *Oreolalax* (*sterlingae* n. sp.) (congeneric comparisons) 213–222; *Microcaecilia* (*savagei* n. sp.) (congeneric comparisons), *Epicrionops niger* (museum specimens) 223–231; *Hypopygus* (*benoneae* n. sp.) 232–237; *Leporinus* (*moralesi, parvulus* n. sp.) 238–247; *Aspidoscelis* (*arizonae, pai, inornata llanuras*) 366–377; *Tatia* (*carolae* n. sp., *marthae* n. sp.) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) 403–414; *Platyberyx* (6 spp.), *Caristius* (4 spp.) 415–434; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) 441–453; *Characidium* (*amaila* n. sp.) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) 464–469; *Arapaima* (*leptosoma* n. sp.) 470–476; *Doumea* (*skeltoni* n. sp., congeners) 577–581; *Acrobrycon* (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) 612–622; *Percina peltata* 623–626; *Cyphocharax* (*aninha* n. sp.) 627–633; *Listrura* (*depinnai* n. sp.) 641–646; *Centrophryne spinulosa* (M/F) 666–669; *Chrosomus* sp. cf. *saylori* (comparison to *C. saylori*) 750–756.

MESOPELAGIC, Caristiidae (family revision; descriptions; distribution maps; spp. key) 415–434; *Centrophryne spinulosa* (sexual parasitism) 666–669.

METHODS, *Macrochelys temminckii* (survivorship; Cormack-Jolly-Seber Model) 58–63; *Montivipera raddei* (radiotelemetry; spatial ecology; minimum convex polygon; home range) 64–71; *Cichlasoma urophthalmus* (regression models; invasive species) 80–96; *Gasterosteus aculeatus* (collagen staining; skeletal ontogeny) 127–131; *Actinemys marmorata* (radiotelemetry; seasonal migration) 142–153; *Psammophylax r. rhombeatus* (radiotelemetry; spatial ecology; minimum convex polygons) 194–200; *Crotalus o. oreganus* (spatial ecology; comparison, MCP vs. FK) 485–492; *Lithobates areolatus* (call survey) 552–561; *Percina peltata* (shape analysis) 622–626; *Pseudacris maculata* (marking methods, survival) 670–675.

MICROSATELLITE DNA, *Gambusia* (*krumholzi*, cf. *clarkhubbsi, speciosa*) (haplotypes) 72–79; *Glyptemys muhlenbergii* (fragmented populations; gene flow) 324–331; *Graptemys caglei* (gene flow; population structure) 723–728.

MIGRATION, *Actinemys marmorata* (seasonal; flow regime influence; radiotelemetry) 142–153; *Leuresthes tenuis* (temporary northern range extension) 729–739.

MINIATURIZATION, *Hypopygus* (*benoneae* n. sp.) (reductive morphological characters) 232–237; *Cyphocharax* (*aninha* n. sp.) (comparison to other small spp.) 627–633.

MITOCHONDRIAL DNA, *Oxygaster* (placement w/in Cyprinidae; combined data matrix) 13–22; *Gambusia* (*krumholzi*, cf. *clarkhubbsi, speciosa*) (population haplotypes) 72–79; *Oreolalax* (*sterlingae* n. sp.) phylogenetic analysis) 213–222; *Hoplias microlepis* (haplotypes; population divergence) 312–323; *Oryzias* (*asinua* n. sp., *wolasi* n. sp., *woworae* sp. gr.) (maximum parsimony, Bayesian inference trees) 403–414.

MORPHOLOGY, *Incilius* (*majordomus* n. sp.) (diagnosis; description; pseudotypanum; coloration; skin; photo./fig.), (congeneric comparisons) 8–12; *Amioides* (diagnosis; comparisons to *Cheilodipterus* and *Synagrops*; table), (*polyacanthus* n. comb.) (diagnosis; redescription; coloration; head pores; osteology; photos.) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) (diagnosis; description; coloration; pseudotypanum; morphometrics; figs.) 31–37; *Arapaima agassizii* (diagnosis; redescription; osteology; original figure, reconstruction) 38–51; *Characidium* (*papachibe* n. sp.) (diagnosis; description; morphometrics; coloration; fig.) 52–57; *Gambusia* (cf. *clarkhubbsi, krumholzi*) (morphological differences) 72–79; *Moxostoma duquesnei* (larval stages; morphometrics; myomeres; figs.) 121–126; *Dendrophidion dendrophis, Mastigodryas bifossatus* (caudal vertebrae, muscles) 132–141; *Actinemys marmorata* (flow regime/body size; morphometrics) 142–153; *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) (dewlap coloration) 201–205; *Oreolalax* (*sterlingae* n. sp.) (diagnosis; description; sexual dimorphism; morphometrics; figs.) 213–222; *Microcaecilia* (*savagei* n. sp.) (diagnosis; description; coloration; figs.; generic measurements), *Epicrionops niger* (redescription; coloration; figs.; measurements, museum specimens) 223–231; *Hypopygus* (*benoneae* n. sp.) (diagnosis; description; morphometrics; coloration; figs.) 232–237; *Leporinus* (*moralesi*) (redescription), (*parvulus* n. sp.) (diagnosis; description), (morphometrics; jaw osteology; coloration; figs.) 238–247; *Hoplias microlepis* (population divergence; environmental factors) 312–323; *Ambystoma maculatum* (sexual dimorphism; visceral organ mass) 338–345; *Aspidoscelis* (*arizonae, pai, inornata llanuras*) (meristics; coloration; scalation) 366–377; *Lamna* (*ditropis, nasus*) (chondrocranium; figs.) 378–389; *Tatia* (*carolae* n. sp., *marthae* n. sp.) (diagnosis; description; coloration, morphometrics; figs.) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (differential diagnoses; descriptions; coloration; mtDNA; meristics/morphometrics; figs.) 403–414; *Platyberyx* (6 spp.), *Caristius* (4 spp.) (generic description), (diagnoses; descriptions; meristics; gill rakers; photos.), (key to spp.) 415–434; *Parotocinclus* (*jequi* n. sp.) (diagnosis; description; sexual dimorphism; coloration; figs.) 435–440; *Typhlobelus* (generic redescription; key to spp.; osteology; synapomorphies), (*auriculatus* n. sp.) (diagnosis; description; morphometrics; coloration; SEM photos.; sexual dimorphism) 441–453; *Characidium* (*amaila* n. sp.) (diagnosis; description; morphometrics; coloration; osteology; swim bladder; figs.) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) (diagnosis; description; coloration; osteology; morphometrics; figs.) 464–469; *Arapaima* (*leptosoma* n. sp.) (diagnosis; description; coloration; morphometrics), (comparisons to nominal spp.) 470–476; *Doumea* (*skeltoni* n. sp.) (diagnosis; description; coloration; morphometrics; fig.) 577–581; †*Lophionotus* (generic redescription), (†*chinleana* n. sp., †*kanabensis* n. comb.) (diagnoses; descriptions; figs.) 582–603; *Acrobrycon* (generic redescription; key to spp.), (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) (diagnoses; descriptions; coloration; morphometrics; secondary sexual dimorphism; figs.) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) (diagnosis; description; coloration; dentition; morphometrics; osteology; figs.) 612–621; *Percina peltata* (meristics; distinctness) 622–626; *Cyphocharax* (*aninha* n. sp.) (diagnosis; description; coloration; morphometrics; osteology; figs.), (congener comparisons) 627–633; *Trachycephalus* (*helioi* n. sp.) (holotype diagnosis, description, coloration, figs.; morphometric variation; single vocal sac) 634–640; *Listrura* (*depinnai* n. sp.) (diagnosis, description, coloration, morphometrics; fig., x-ray) 641–646; *Centrophryne spinulosa* (M/F description; figs.) 666–669; *Rana sylvatica* (metamorphosis & effects of canopy cover) 717–722; *Aplodinotus grunniens* (variation, effects of environmental factors; allometry) 740–749; *Chrosomus* sp. cf. *saylori* (ontogeny; meristics), (comparison to *C. saylori*) 750–756.

MORTALITY RATE, *Desmognathus* (*monticola, ocoee, quadramaculatus*) (body size, reproduction) 262–267; *Podocnemis unifilis* (hatching success; survival) 303–311; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (increased salinity; lethal/sublethal concentrations) 562–566; *Pseudacris maculata* (marking methods & survival; probability models) 670–675; *Lithobates* (*catesbeianus, clamitans*) (predator–prey interactions, sublethal pesticides) 691–698.

MOVEMENT [see also MIGRATION], *Montivipera raddei* (spatial ecology; habitat preference; modified landscape) 64–71; *Psammophylax r. rhombeatus* (spatial ecology; M/F home ranges) 194–200; *Crotalus o. oreganus* (spatial ecology; M/F home ranges; environmental factors) 485–492; *Heterodon platirhinos* (road crossing; avoidance) 507–511; *Desmognathus ocoee* (jumping; kinematics; defense mechanism) 512–516; *Etheostoma okaloosae* (mark/recapture; age/longevity) 653–659; *Bothrops atrox* (habitat selection) 684–690.

MUSEUMS, *Arapaima agassizii* (holotype loss; osteological drawings) 38–51; *Gambusia* (cf. *clarkhubbsi, krumholzi*) (morphological differences) 72–79; *Epicrionops niger* (table, specimen measurements) 223–231.

NATURAL HISTORY [see also LIFE HISTORY], *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. (trophic interactions; helminth parasite transmission) 103–110; *Dendrophidion dendrophis,*

Mastigodryas bifossatus (pseudoautotomy) 132–141; *Etheostoma luteovinctum* (male mating choice; visual cues) 154–159; *Liolaemus chiliensis* (distress call) 206–212; *Lithobates clamitans* (tadpoles, abiotic/biotic environmental factors) 248–253; *Ambystoma* (*talpoideum, maculatum*) (intraguild predation; cannibalism) 284–291; *Podocnemis unifilis* (nests, environmental factors; hatching success; sex ratio) 303–311; *Ambystoma maculatum* (breeding; sexual dimorphism) 338–345; *Typhlobelus* (*auriculatus* n. sp.) (psammophilic; sexual dimorphism) 441–453; *Graptemys geographica* (diet, sexual differences) 477–484; *Crotalus o. oreganus* (spatial ecology; environmental factors; movement, sexual differences) 485–492; *Thalassoma bifasciatum* (larval settlement; spawning) 493–498; Crocodylia (long-distance signaling) 517–526; *Anaxyrus* (=*Bufo*) *fowleri* (breeding sites, sex ratio, population size) 647–652; *Etheostoma okaloosae* (movement; longetivity) 653–659; *Sladenia shaefersi* (*in-situ* observations; photos.) 660–665; *Plethodon cinereus* (assortative mating) 676–683; *Bothrops atrox* (habitat selection) 684–690; *Pelobates syriacus* (reproductive strategies) 699–706; *Leuresthes tenuis* (northern habitat expansion) 729–739.

NESTS/NESTING, *Podocnemis* (*expansa, unifilis, vogli*) (site choice; illegal harvesting) 111–120; *Alligator mississippiensis* (habitat quality, size) 268–276; *Podocnemis unifilis* (environmental factors; hatching success; survival; clutch size; sex ratio) 303–311.

NOMENCLATURE, *Incilius* (*majordomus* n. sp.) (history Neotropical bufonids) 8–12; Oxygastrinae (revision; subfamily synonyms) 13–22; *Amioides* (*polyacanthus* n. comb.) (validity) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) 31–37; *Arapaima agassizii* (validity) 38–51; *Characidium* (*papachibe* n. sp.) 52–57; *Gambusia* (cf. *clarkhubbsi*) (junior synonym) 72–79; *Oreolalax* (*sterlingae* n. sp.) 213–222; *Microcaecilia* (*savagei* n. sp.) 223–231; *Hypopygus* (*benoneae* n. sp.) 232–237; *Leporinus moralesi* (junior synonym; lectotype designation), (*parvulus* n. sp.) 238–247; *Aspidoscelis arizonae* (morphometrics; diagnosis; validity) 366–377; *Tatia* (*carolae* n. sp., *marthae* n. sp.), *musaica* (generic placement) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) 403–414; *Platyberyx* (3 n. spp.), *Caristius* (2 n. spp.) 415–434; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) 441–453; *Characidium* (*amaila* n. sp.) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) 464–469; *Arapaima* (*leptosoma* n. sp.) 470–476; *Doumea* (*skeltoni* n. sp.) 577–581; †*Lophionotus* (*chinleana* n. sp., *kanabensis* n. comb.) 582–603; *Acrobrycon* (*starnesi* n. sp., *ortii* n. sp.), (*tarijae* placed in synonymy) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) 612–621; *Cyphocharax* (*aninha* n. sp.) 627–633; *Trachycephalus* (*helioi* n. sp.) 634–640; *Listrura* (*depinnai* n. sp.) 641–646.

NUCLEAR DNA, Oxygastrinae (cyprinid phylogenetic position; mtDNA, combined data matrix) 8–12.

OBITUARIES, R. H. Backus 173–176; J. B. Graham 346–350; N. V. Parin 768–780.

ONTOGENY, *Moxostoma duquesnei* (developmental stages; morphometrics; figs.) 121–126; *Gasterosteus aculeatus* (collagen straining method) 127–131; *Chrosomus* sp. cf. *saylori* (comparison to congeners; morphologic variation) 750–756.

OSTEOLOGY, *Amioides* (*polyacanthus* n. comb.) (x-rays, full body) 23–30; *Arapaima agassizii* (original drawings; redescription; congener comparison) 38–51; *Dendrophidion dendrophis, Mastigodryas bifossatus* (caudal vertebrae; pseudoautotomy) 132–141; *Leporinus* (*moralesi, parvulus* n. sp.) (jaws; figs.) 238–247; *Lamna* (*ditropis, nasus*) (chondrocranium; figs.) 378–389; *Typhlobelus* (*auriculatus* n. sp.) (cranium, figs.) 441–453; *Characidium* (*amaila* n. sp.) (neurocranium; vertebrae; figs.) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) (upper & lower jaws; fig.) 464–469; †*Lophionotus* (*chinleana* n. sp., *kanabensis* n. comb.) (descriptions; figs.) 582–603; *Hyphessobrycon* (*kayabi* n. sp.) (anal fin; dentition; figs.) 612–621; *Cyphocharax* (*aninha* n. sp.) (infraorbital series; congener comparisons; fig.) 627–633.

OTOLITH, *Moxostoma* (*duquesnei, erythrurum*) (age; growth) 390–395; *Thalassoma bifasciatum* (planktonic phase; fig.) 493–498.

PAEDOMORPHISM, *Ambystoma talpoideum* (predation) 284–291.

PALEONTOLOGY, †*Lophionotus* (*chinleana* n. sp.) (U. Triassic, Chinle Fm.), (*kanabensis* n. comb.) (L. Jurassic, Moenave Fm.) 582–603; *Cyphocharax* (*aninha* n. sp.) (miniaturization characters) 627–633.

PARASITES/PARASITISM, *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. (helminthes, transmission; tropic interactions) 103–110; *Centrophryne spinulosa* (sexual, 1st record) 666–669.

PATHOGENS/PATHOLOGY, *Sistrurus c. catenatus* (*Chrysosporium* fungus; hematology; plasma biochemistry; heavy metals) 97–102.

PESTICIDES, *Lithobates* (*catesbeianus, clamitans*) (sublethal effects, predator–prey interactions) 691–698.

PHYLOGENETIC ANALYSIS, *Oxygaster*, Oxygastrinae (mtDNA, nuclear DNA; maximum likelihood tree; cyprinid placement) 13–22; *Oreolalax* (*sterlingae* n. sp.) (mtDNA; maximum parsimony tree; species placement) 213–222; *Oryzias* (*asinua* n. sp., *wolasi* n. sp., *woworae* sp. gr.) (mtDNA) 403–414; †Semionotidae, Ginglymodi (morphological data) 582–603.

PHYLOGENY, *Oxygaster* (cyprinid placement; historical genetic analysis) 13–22; *Oreolalax* (*sterlingae* n. sp.) (generic molecular analysis, morphological comparisons) 213–222; *Typhlobelus* (rediagnosis) 441–453; *Listrura* (*depinnai* n. sp.) (inclusion in Glanapteryginae) 641–646.

PHYSIOLOGY, *Sistrurus c. catenatus* (hematology, plasma biochemistry, heavy metals, fungal pathogens) 97–102; *Actinemys marmorata* (body condition; flow regime influence) 142–153; *Bufo melanostictus* (plasma sex steroids) 332–337; *Ambystoma maculatum* (sexual dimorphism; hematology) 338–345; *Rhinella marina* (corticosterone levels; stress response) 539–542; anuran larvae (7 spp.) (salinity level tolerance) 543–551; *Crotalus lepidus morulus* (infraspecific venom variation) 707–716.

PIGMENTATION [see also COLORATION], *Moxostoma duquesnei* (larval development) 121–126; *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) (dewlap color variation; pterins/carotenoids) 201–205; *Leporinus* (*moralesi, parvulus* n. sp.) (diagnostic patterns; figs.) 238–247.

POPULATIONS, *Macrochelys temminckii* (commercial harvesting; recovery; survivorship) 58–63; *Gambusia krumholzi* (restricted distribution; conservation status) 72–79; *Cichlasoma urophthalmus* (invasive species, estuarine Everglades; effect on native fish) 80–96; *Alligator mississippiensis* (marginality, reproductive allometry) 268–276; *Hoplias microlepis* (morphologic/genetic divergence; environment factors; impoundments) 312–323; *Glyptemys muhlenbergii* (fragmented; gene flow; microsatellite DNA; conservation) 324–331; *Percina peltata* (isolated; distinct morphometrics) 622–626; *Anaxyrus* (=*Bufo*) *fowleri* (breeding sites; sex ratio) 647–652; *Graptemys caglei* (gene flow; environmental factors; microsatellite DNA) 723–728; *Leuresthes tenuis* (temporary northern range expansion) 729–739.

PREDATION, *Lampropholis delicata* (predator escapement; early hatching; sensory detection, survival mechanism) 160–165; *Gambusia affinis* (effect on *Lithobates clamitans* tadpoles) 248–253; *Micropterus salmoides* ("dangerous prey" hypothesis; spine avoidance) 277–283; *Ambystoma* (*talpoideum, maculatum*) (intraguild; cannibalism) 284–291; *Lithobates* (*catesbeianus, clamitans*) (sublethal effects of pesticides) 691–698.

PSEUDOAUTOTOMY, *Dendrophidion dendrophis, Mastigodryas bifossatus* (caudal vertebrae; muscles) 132–141.

PSEUDOTYMPANUM, *Incilius* (*majordomus* n. sp.) 8–12; *Hemigrammocharax* (*rubensteini* n. sp.) (first report for Distichodontidae) 31–37; *Typhlobelus* (*auriculatus* n. sp.) (modified; unique; SEM figs.) 441–453.

REPRODUCTION, *Alligator mississippiensis* (allometry; egg/clutch/nest size; habitat quality) 268–276; *Podocnemis unifilis* (nesting site; hatching success) 303–311; *Bufo melanostictus* (behavior; plasma sex steroids; "associated reproductive pattern") 332–337; *Ambystoma maculatum* (sexual dimorphism, breeding; hematology) 338–345; *Crotalus o. oreganus* (movement, mating season, body size) 485–492; *Thalassoma bifasciatum* (spawning) 493–498; *Fundulus heteroclitus* (spawning in lab environment; egg production) 527–538; *Sladenia syriacus* (*in-situ* observations; photos.) 660–665; *Centrophryne spinulosa* (1st record, sexual parasitism; figs.) 666–669; *Plethodon cinereus* (assortative mating; scent/diet/territoriality factors) 676–683; *Pelobates syriacus* (strategies) 699–706.

SALINITY, anuran larvae (7 spp.) (concentration increase; tolerance; survival) 543–551; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (concentration increase; lethal/sublethal effects) 562–566.

SCALATION, *Aspidoscelis* (*arizonae, pai, inornata llanuras*) (diagnostic reliability) 366–377.

SENSORY, *Lampropholis delicata* (hatchling vibration detection; predator escapement; survival mechanism) 160–165.

SETTLEMENT, *Thalassoma bifasciatum* (larval stage duration) 493–498.

SEX, *Pelobates syriacus* (reproductive strategies) 699–706; *Aplodinotus grunniens* (effect on morphologic variation) 740–749.

SEX RATIO, *Macrochelys temminckii* (female bias) 58–63; *Podocnemis* (*expansa, unifilis, vogli*) (effect of harvesting) 111–120; *Podocnemis unifilis* (nesting site selection; hatching success; environmental factors) 303–311; *Anaxyrus* (= *Bufo*) *fowleri* (breeding site population) 647–652.

SEXUAL BEHAVIOR, *Etheostoma luteovinctum* (male mating preferences; visual cues) 154–159; *Bufo melanostictus* (plasma sex steroids) 332–337; *Graptemys geographica* (M/F diet differences) 477–484; *Centrophryne spinulosa* (male parasitism) 666–669; *Plethodon cinereus* (assortative mating; mechanisms) 676–683.

SEXUAL DICHROMATISM, *Incilius* (*majordomus* n. sp.) 8–12; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (description; photos.) 403–414.

SEXUAL DIMORPHISM, *Incilius* (*majordomus* n. sp.) (coloration) 8–12; *Characidium* (*papachibe* n. sp.) (absence) 52–57; *Oreolalax* (*sterlingae* n. sp.) 213–222; *Ambystoma maculatum* (visceral organ mass; hematology; breeding) 338–345; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) 441–453; *Acrobrycon* (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) (fin-ray bony hooks) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) (anal fin; fig.) 612–621; *Sladenia shaefersi* (size; photos.) 660–665; *Pelobates syriacus* (size) 699–706.

SEXUAL PARASITISM, *Centrophryne spinulosa* (1st record; figs.) 666–669.

SIZE, *Podocnemis* (*expansa, unifilis, vogli*) (carapace/plastron; harvesting effect) 111–120; *Actinemys marmorata* (body size; morphometrics; flow regime influence) 142–153; *Desmognathus* (*monticola, ocoee, quadramaculatus*) (body size/fecundity) 262–267; *Alligator mississippiensis* (reproductive allometry) 268–276; *Crotalus o. oreganus* (body size, movement, home range, mating season) 485–492; *Fundulus heteroclitus* (gonads wrt latitude) 527–538; *Anaxyrus* (= *Bufo*) *fowleri* (breeding site population, abundance over time) 647–652; *Rana sylvatica* (effects of canopy cover) 717–722; *Aplodinotus grunniens* (morphologic variation; environmental factors) 740–749.

SKIN, *Incilius* (*majordomus* n. sp.) (tubercle absence; comparison to *I. peripatetes*; fig.) 8–12.

SOUND, *Liolaemus chiliensis* (vocalization) 206–212; Crocodylia (long-distance signaling) 517–526.

SPATIAL ECOLOGY, *Montivipera raddei* (movement between habitats) 64–71; *Psammophylax r. rhombeatus* (movement, sexual difference) 194–200; *Ambystoma* (*laterale-jeffersonianum* complex) (habitat selection; chemical cues) 254–261; *Glyptemys muhlenbergii* (fragmented populations; gene flow) 324–331; *Crotalus o. oreganus* (home range; body size; environmental factors) 485–492; *Desmognathus* (*folkertsi, quadramaculatus*) (trophic interaction; competition-predation hypothesis) 499–506; *Heterodon platirhinos* (road-crossing mortality rate) 507–511; *Bothrops atrox* (habitat selection, movement) 684–690; *Leuresthes tenuis* (northern habitat expansion) 729–739.

SURVIVAL/SURVIVORSHIP, *Macrochelys temminckii* (commercial harvesting; population recovery) 58–63; *Montivipera raddei* (human-modified landscape; spatial ecology) 64–71; *Lampropholis delicata* (early hatching; predator detection; sensory mechanism) 160–165; *Lithobates clamitans* (tadpoles, abiotic/biotic environmental factors) 248–253; *Desmognathus* (survival to 1st reproduction) 262–267; *Podocnemis unifilis* (nests; hatching success; environmental factors) 303–311; anuran larvae (7 spp.) (salinity level tolerance) 543–551; *Lithobates berlandieri, Incilius nebulifer, Acris crepitans* (salinity, lethal/sublethal effects) 562–566; *Pseudacris maculata* (marking methods) 670–675.

SWIM BLADDER, *Characidium* (*amaila* n. sp.) (fig.) 454–463.

SYSTEMATICS, *Incilius* (*majordomus* n. sp.) (Neotropical Bufonidae placement) 8–12; *Oxygaster* (family placement) 13–22; *Amioides* (*polyacanthus* n. comb.) (family placement; generic validity) 23–30; *Hemigrammocharax-Nannocharax* clade (pseudotypanum significance) 31–37; *Arapaima agassizii* (species validity) 38–51; *Characidium* (*papachibe* n. sp.) (comparison to psammophile congeners) 52–57; *Microcaecilia* (*savagei* n. sp.) 223–231; *Hypopygus* (*benoneae* n. sp.) (synapomorphies) 232–237; *Leporinus* (*moralesi, uatumaensis, muelleri, parvulus* n. sp.) 238–247; *Tatia* (*carolae* n. sp., *marthae* n. sp.; *musaica*) (generic placement) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (placement in *woworae* sp. gr.) 403–414; *Platyberyx* (6 spp.), *Caristius* (4 spp.) (family revision) 415–434; *Parotocinclus* (*jequi* n. sp.) 435–440; *Typhlobelus* (*auriculatus* n. sp.) 441–453; *Characidium* (*amaila* n. sp.) (phylogenetic placement w/in genus) 454–463; *Tomistoma schleglii* (position w/in Crocodylia) 517–526; Ginglymodi, †Semionotidae (phylogenetic analysis; synapomorphies) 582–603; *Acrobrycon* (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) (PCA; synonymy of *tarijae*) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) (congeneric relationship) 612–621; *Cyphocharax* (*aninha* n. sp.) (morphological comparisons to congeners) 627–633; *Trachycephalus* (*helioi* n. sp.) (comparisons to congeners) 634–640; *Listrura* (*depinnai* n. sp.) (placement in Glanapteryginae; putative autapomorphies) 641–646.

TADPOLES, *Lithobates clamitans* (abiotic/biotic environmental factors, survivorship; prey of *Gambusia affinis*) 248–253; *Lithobates* (*catesbeianus, clamitans*) (sublethal pesticides, predator–prey interactions) 691–698.

TAXONOMY, *Incilius* (*majordomus* n. sp.) (description; fig.) 8–12; Oxygastrinae (status; genera; synonymized subfamilies) 13–22; *Amioides* (*polyacanthus* n. comb.) (redescription; validity) 23–30; *Hemigrammocharax* (*rubensteini* n. sp.) (description; figs.) 31–37; *Arapaima agassizii* (redescription; figs.) 38–51; *Characidium* (*papachibe* n. sp.) (description; fig.) 52–57; *Gambusia* (*krumholzi*, cf. *clarkhubbsi, speciosa*) (haplotypes; species status; hybridization) 72–79; *Oreolalax* (*sterlingae* n. sp.) (description; figs.) 213–222; *Microcaecilia* (*savagei* n. sp.), *Epicrionops niger* (description; figs.) 223–231; *Hypopygus* (*benoneae* n. sp.) (description; figs.) 232–237; *Leporinus moralesi* (redescription), (*parvulus* n. sp.) (description) 238–247; *Aspidoscelis arizonae* (species status validity; diagnostics) 366–377; *Tatia* (*carolae* n. sp., *marthae* n. sp.) (descriptions; photos.), *musaica* (generic placement) 396–402; *Oryzias* (*asinua* n. sp., *wolasi* n. sp.) (descriptions; photos.) 403–414; *Platyberyx, Caristius* (family revision; spp. key), (generic and species descriptions) 415–434; *Parotocinclus* (*jequi* n. sp.) (description; figs.) 435–440; *Typhlobelus* (generic redescription; synapomorphies), (*auriculatus* n. sp.) (description, figs.) 441–453; *Characidium* (*amaila* n. sp.) (description, figs.; plesiomorphies) 454–463; *Hypomasticus* (*lineomaculatus* n. sp.) (description, figs.; relation to *despaxi*) 464–469; *Arapaima* (*leptosoma* n. sp.) (description, figs.) 470–476; *Doumea* (*skeltoni* n. sp.) (description, fig.) 577–581; †*Lophionotus* (redescription), (*chinleana* n. sp., *kanabensis* n. comb.) (descriptions; figs.) 582–603; *Acrobrycon* (redescription), (*ipanquianus, starnesi* n. sp., *ortii* n. sp.) (descriptions, figs.) 604–611; *Hyphessobrycon* (*kayabi* n. sp.) 612–621; *Percina peltata* (isolated population; distinct morphometrics) 622–626; *Cyphocharax* (*aninha* n. sp.) (description; fig.) 627–6333; *Trachycephalus* (*helioi* n. sp.) (description; figs.) 634–640; *Listrura* (*depinnai* n. sp.) (description; figs.) 641–646.

TEMPERATURE-DEPENDENT SEX DETERMINATION (TSD), *Podocnemis unifilis* (sex ratio; environmental factors) 303–311.

TROPHIC INTERACTIONS, *Perccottus glenii, Natrix* (*natrix, tessellata*), *Pelophylax* spp. (feeding; parasite transmission) 103–110; *Lithobates clamitans* (tadpoles, abiotic/biotic environmental factors) 248–253; *Ambystoma* (*talpoideum, maculatum*) (intraguild predation) 284–291; *Desmognathus* (*folkertsi, quadramaculatus*) (competition-predation hypothesis) 499–506; *Chrosomus* sp. cf. *saylori* (diet preference; comparison to congeners) 750–756.

UROTOMY, *Dendrophidion dendrophis, Mastigodryas bifossatus* (tail morphology; breakage frequency) 132–141.

VARIATION, *Anolis jubar* ssp. (*oriens, maisiensis, albertschwartzi*) (dewlap coloration; pterins/carotenoids) 201–205; *Aspidoscelis arizonae* (coloration; morphological diagnosis) 366–377; *Characidium* (*amaila* n. sp.) (PCA, morphometrics) 454–463;

Fundulus heteroclitus (egg production wrt latitude) 527–538; anuran larvae (7 spp.) (salinity level tolerance; survival) 543–551; *Percina peltata* (morphometrics; isolated population) 622–626; *Trachycephalus* (*helioi* n. sp.) (morphometrics, males only) 634–640; *Crotalus lepidus morulus* (infraspecific venom analysis) 707–716; *Aplodinotus grunniens* (morphologic; environmental factors) 740–749.

VENOM, *Crotalus lepidus morulus* (infraspecific variation) 707–716.

VOLUME CONTENTS

Copeia 2013, Nos. 1–4

Abrahão, Carlos R. (*see* **de Fraga, Rafael**)

Acord, Megan A., Carl D. Anthony, and Cari-Ann M. Hickerson. Assortative Mating in a Polymorphic Salamander 676

Aghasyan, Aram L. (*see* **Ettling, Jeffery A.**)

Aghasyan, Levon A. (*see* **Ettling, Jeffery A.**)

Agorastos, Stergiani (*see* **Aguirre, Windsor E.**)

Agudelo-Zamora, Henry (*see* **Lujan, Nathan K.**)

Aguirre, Windsor E., Virginia R. Shervette, Ronald Navarrete, Paola Calle, and Stergiani Agorastos. Morphological and Genetic Divergence of *Hoplias microlepis* (Characiformes: Erythrinidae) in Rivers and Artificial Impoundments of Western Ecuador 312

Alfonso, Yasel U., Humberto J. Morris, Adrián Gutiérrez, Lourdes Rodríguez-Schettino, Dennis Denis, and John E. Steffen. Dewlap Color Variation Based on Pterin and Carotenoid Pigments in Three Subspecies of *Anolis jubar* of the Cuban Southern Coast 201

Allen, David (*see* **Belasen, Anat**)

Allender, Matthew C., Michael J. Dreslik, Daniel B. Wylie, Sarah J. Wylie, John W. Scott, and Christopher A. Phillips. Ongoing Health Assessment and Prevalence of *Chrysosporium* in the Eastern Massasauga (*Sistrurus catenatus catenatus*) 97

Anderson, Thomas L., Cy L. Mott, Todd D. Levine, and Howard H. Whiteman. Life Cycle Complexity Influences Intraguild Predation and Cannibalism in Pond Communities 284

Anderson, William D., Jr. Field Guide to Fishes of the Chesapeake Bay. By Edward O. Murdy and John A. Musick, Illustrated by Val Kells 784

Anthony, Carl D. (*see* **Acord, Megan A.**)

Arcila, Dahiana, Richard P. Vari, and Naércio A. Menezes. Revision of the Neotropical Genus *Acrobrycon* (Ostariophysi: Characiformes: Characidae) with Description of Two New Species 604

Babitzke, Joel B. (*see* **Ward, Rocky**)

Backus, Edward H., and Bruce B. Collette. Richard Haven Backus (1922–2012) 173

Bailey, Larissa L. (*see* **Swanson, Jennifer E.**)

Baird, Troy A., Kathryn A. Dickson, and Nicholas C. Wegner. Jeffrey Brent Graham (1941–2011) 346

Baker, Sherri (*see* **Echelle, Anthony A.**)

Băncilă, Raluca I. (*see* **Cogălniceanu, Dan**)

Barreto, Guillermo R. (*see* **Peñaloza, Claudia L.**)

Bauer, Aaron M. (*see* **Cottone, Amanda M.**)

Beckman, Daniel W., and David T. Howlett. Otolith Annulus Formation and Growth of Two Redhorse Suckers (*Moxostoma*: Catastomidae) 390

Belasen, Anat, Erin Burkett, Allison Injaian, Kevin Li, David Allen, and Ivette Perfecto. Effect of Sub-Canopy on Habitat Selection in the Blue-spotted Salamander (*Ambystoma laterale-jeffersonianum* unisexual complex) 254

Bell, Michael A. (*see* **Lohman, Brian K.**)

Benard, Michael F. (*see* **Boes, Matthew W.**)

Birindelli, José L. O., Heraldo A. Britski, and Flávio C. T. Lima. New Species of *Leporinus* from the Rio Tapajós Basin, Brazil, and Redescription of *L. moralesi* (Characiformes: Anostomidae) 238

Birindelli, José L. O., Luiz A. W. Peixoto, Wolmar B. Wosiacki, and Heraldo A. Britski. New Species of *Hypomasticus* Borodin, 1929 (Characiformes: Anostomidae) from Tributaries of the Lower Rio Amazonas, Brazil 464

Biscoito, Manuel (*see* **Vieira, Sofia**)

Blouin-Demers, Gabriel (*see* **Robson, Laura E.**)

Boes, Matthew W., and Michael F. Benard. Carry-Over Effects in Nature: Effects of Canopy Cover and Individual Pond on Size, Shape, and Locomotor Performance of Metamorphosing Wood Frogs 717

Böhme, Wolfgang (*see* **Nguyen, Truong Q.**)

Bondi, Cheryl A., and Sharyn B. Marks. Differences in Flow Regime Influence the Seasonal Migrations, Body Size, and Body Condition of Western Pond Turtles (*Actinemys marmorata*) that Inhabit Perennial and Intermittent Riverine Sites in Northern California 142

Booth, Pieter N. (*see* **Lujan, Nathan K.**)

Borja, Miguel, David Lazcano, Gerardo Martínez-Romero, Jesús Morlett, Elda Sánchez, Ana C. Cepeda-Nieto, Yolanda Garza-García, and Alejandro Zugasti-Cruz. Intra-specific Variation in the Protein Composition and Proteolytic Activity of Venom of *Crotalus lepidus morulus* from the Northeast of Mexico 707

Bosker, Thijs, Kelly R. Munkittrick, Diane E. Nacci, and Deborah L. MacLatchy. Laboratory Spawning Patterns of Mummichogs, *Fundulus heteroclitus* (Cyprinodontiformes: Fundulidae) 527

Braun, Bruna Koech (*see* **Lehmann A., Pablo**)

Britski, Heraldo A. (*see* **Birindelli, José L. O.**)

Brown, Mary E., and Susan C. Walls. Variation in Salinity Tolerance among Larval Anurans: Implications for Community Composition and the Spread of an Invasive, Non-native Species 543

Bruce, Richard C. Size-Mediated Tradeoffs in Life-History Traits in Dusky Salamanders 262

Bunt, Christopher M., Thom Heiman, and Nicholas E. Mandrak. Ontogeny of Larval and Juvenile Black Redhorse (*Moxostoma duquesnei*) 121

Burkett, Erin (*see* **Belasen, Anat**)

Burner, Allison B. (*see* **Smith, Geoffrey R.**)

Calle, Paola (*see* **Aguirre, Windsor E.**)

Camp, Carlos D., Jessica A. Wooten, Caitlin M. Corbet, Eden A. Dulka, Joshua A. Mitchem, and Thomas J. Krieger. Ecological Interactions between Two Broadly Sympatric, Cryptic Species of Dusky Salamander (Genus *Desmognathus*) 499

Caruso, John H. (*see* **Pietsch, Theodore W.**)

Cepeda-Nieto, Ana C. (*see* **Borja, Miguel**)

Chakrabarty, Prosanta. Systematics, Biology, and Distribution of the Species of the Oceanic Oarfish Genus *Regalecus* (Teleostei, Lampridiformes, Regalecidae). By Tyson R. Roberts 356

© 2013 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/OT-13-138

Chikhlyaev, Igor V. (*see* **Reshetnikov, Andrey N.**)
Ciccotto, Patrick J. Morphometric Distinctness of an Isolated Population of *Percina peltata* (Shield Darter) on the Atlantic Coastal Plain of Maryland, USA 622
Ciccotto, Patrick J., Jennifer M. Gumm, and Tamra C. Mendelson. Male Association Preference for Conspecifics in the Redband Darter, *Etheostoma luteovinctum* (Teleostei: Percidae) Based on Visual Cues 154
Cogălniceanu, Dan, Paul Székely, Diana Székely, Daniela Roşioru, Raluca I. Băncilă, and Claude Miaud. When Males Are Larger than Females in Ecthotherms: Reproductive Investment in the Eastern Spadefoot Toad *Pelobates syriacus* 699
Collette, Bruce B. Nikolai Vasil'evich Parin (1932–2012) 768
Collette, Bruce B. (*see* **Backus, Edward H.**)
Cooper, Amos (*see* **Murray, Christopher M.**)
Corbet, Caitlin M. (*see* **Camp, Carlos D.**)
Cordes, James E. (*see* **Sullivan, Brian K.**)
Cottone, Amanda M., and Aaron M. Bauer. The Vernal Spatial Ecology and Mating Behaviors of the Rhombic Skaapsteker, *Psammophylax rhombeatus rhombeatus* (Serpentes: Psammophiidae), from the Western Cape, South Africa 194
Cowman, Peter F. Discovery of Australia's Fishes: A History of Australian Ichthyology to 1930. By Brian Saunders 786
Crother, Brian I. (*see* **Murray, Christopher M.**)
Crowder, Larry B. (*see* **Peñaloza, Claudia L.**)
Das, Indraneil. A Contribution to the Herpetology of Northern Pakistan. By R. Masroor 178
Dayer, Christopher B. (*see* **Smith, Geoffrey R.**)
de Fraga, Rafael, William E. Magnusson, Carlos R. Abrahão, Tania Sanaiotti, and Albertina P. Lima. Habitat Selection by *Bothrops atrox* (Serpentes: Viperidae) in Central Amazonia, Brazil 684
de Pinna, Mário C. C., and Jansen Zuanon. The Genus *Typhlobelus*: Monophyly and Taxonomy, with Description of a New Species with a Unique Pseudotympanic Structure (Teleostei: Trichomycteridae) 441
de Santana, Carlos David (*see* **Peixoto, Luiz Antônio Wanderley**)
Delgado, João (*see* **Vieira, Sofia**)
Denis, Dennis (*see* **Alfonso, Yasel U.**)
Dibble, Christopher J. (*see* **Smith, Geoffrey R.**)
Dickson, Kathryn A. (*see* **Baird, Troy A.**)
Dinets, Vladimir. Long-Distance Signaling in Crocodylia 517
Dinkelacker, Stephen A. (*see* **Howey, Christopher A. F.**)
Donnelly, Maureen A. Jay M. Savage 757
Donnelly, Maureen A., and Marvalee H. Wake. A New *Microcaecilia* (Amphibia: Gymnophiona) from Guyana with Comments on *Epicrionops niger* 223
Donnelly, Maureen A. (*see* **Savage, Jay M.**)
Doody, J. Sean, and Phillip Paull. Hitting the Ground Running: Environmentally Cued Hatching in a Lizard 160
Douglas, Marlis R. (*see* **Sullivan, Brian K.**)
Douglas, Michael E. (*see* **Sullivan, Brian K.**)
Dourado, Ângelo C. M., Leonardo Oliveira, and Ana L. C. Prudente. Pseudoautotomy in *Dendrophidion dendrophis* and *Mastigodryas bifossatus* (Serpentes: Colubridae): Tail Morphology and Breakage Frequency 132
Dreslik, Michael J. (*see* **Allender, Matthew C.**)
Dulka, Eden A. (*see* **Camp, Carlos D.**)
Dutra, Guilherme Moreira (*see* **Peixoto, Luiz Antônio Wanderley**)
Easter, Michael (*see* **Murray, Christopher M.**)
Ebert, David A. Global Perspectives on the Biology and Life History of the White Shark. By Michael L. Dormeier (ed.) 351
Echelle, Alice F. (*see* **Echelle, Anthony A.**)
Echelle, Anthony A., Ma. de Lourdes Lozano Vilano, Sherri Baker, Wade D. Wilson, Alice F. Echelle, Gary P. Garrett, and Robert J. Edwards. Conservation Genetics of *Gambusia krumholzi* (Teleostei: Poeciliidae) with Assessment of the Species Status of *G. clarkhubbsi* and Hybridization with *G. speciosa* 72
Edwards, Robert J. (*see* **Echelle, Anthony A.**)
Encarnação, Helena (*see* **Vieira, Sofia**)
Engbrecht, Nathan J. (*see* **Williams, Perry J.**)
Espín, Rodolfo (*see* **Peñaloza, Claudia L.**)
Ettling, Jeffery A., Levon A. Aghasyan, Aram L. Aghasyan, and Patricia G. Parker. Spatial Ecology of Armenian Vipers, *Montivipera raddei*, in a Human-Modified Landscape 64
Fayzulin, Alexandr I. (*see* **Reshetnikov, Andrey N.**)
Fernandes, Luana F. (*see* **Pignati, Marina T.**)
Ferraris, Carl J., Jr., and Richard P. Vari. New Species of *Doumea* (Siluriformes: Amphiliidae) from Coastal River Basins of the Republic of the Congo 577
Ferraris, Carl J., Jr. (*see* **Vari, Richard P.**)
Ferreira, Paulo D. (*see* **Pignati, Marina T.**)
Ferrer, Juliano (*see* **Villa-Verde, Leandro**)
Fine, Michael L. (*see* **Sismour, Edward N.**)
Finkler, Michael S. Sexual Dimorphism in Visceral Organ Mass and Hematology in Spotted Salamanders, *Ambystoma maculatum*, before and after Breeding 338
Fisher, Charles R. (*see* **Pietsch, Theodore W.**)
Fisher, Kaitlin J., Kevin J. Guilfoyle, and Kent A. Hatch. Stress Induced by Toe-Clipping in Cane Toads (*Rhinella marina*) 539
Fraser, Thomas H. Family Status of *Amioides* and the Senior Synonym of *Coranthus* (Percomorpha: Apogonidae) 23
Funk, W. Chris (*see* **Swanson, Jennifer E.**)
Garrett, Gary P. (*see* **Echelle, Anthony A.**)
Garza-García, Yolanda (*see* **Borja, Miguel**)
Gibbs, James P. (*see* **Shoemaker, Kevin T.**)
Gibson, Sarah Z. Biodiversity and Evolutionary History of †*Lophionotus* (Neopterygii: †Semionotiformes) from the Western United States 582
Gordo, Marcelo (*see* **Nunes, Ivan**)
Gramapurohit, Narahari P., and Rajkumar S. Radder. Changes in Plasma Sex Steroids in Relation to Reproductive Behavior of the Toad, *Bufo melanostictus* (Schn.) 332
Green, David M. Sex Ratio and Breeding Population Size in Fowler's Toad, *Anaxyrus* (=*Bufo*) *fowleri* 647
Gresens, Susan E. (*see* **Richards-Dimitrie, Teal**)
Guilfoyle, Kevin J. (*see* **Fisher, Kaitlin J.**)
Gumm, Jennifer M. (*see* **Ciccotto, Patrick J.**)
Gutiérrez, Adrián (*see* **Alfonso, Yasel U.**)
Hadiaty, Renny K. (*see* **Parenti, Lynne R.**)
Hanlon, Shane M., and Rick Relyea. Sublethal Effects of Pesticides on Predator–Prey Interactions in Amphibians 691
Harrison, Elizabeth, Jerome J. Lorenz, and Joel C. Trexler. Per Capita Effects of Non-native Mayan Cichlids (*Cichlasoma urophthalmus*; Gunther) on Native Fish in the Estuarine Southern Everglades 80

Hatch, Kent A. (*see* **Fisher, Kaitlin J.**)
Heiman, Thom (*see* **Bunt, Christopher M.**)
Herder, Fabian (*see* **Parenti, Lynne R.**)
Hernández, Omar (*see* **Peñaloza, Claudia L.**)
Hickerson, Cari-Ann M. (*see* **Acord, Megan A.**)
Hieb, Kathryn A. (*see* **Martin, Karen L. M.**)
Holt, Daniel E., Howard L. Jelks, and Frank Jordan. Movement and Longevity of Imperiled Okaloosa Darters (*Etheostoma okaloosae*) 653
Howey, Christopher A. F., and Stephen A. Dinkelacker. Characteristics of a Historically Harvested Alligator Snapping Turtle (*Macrochelys temminckii*) Population 58
Howlett, David T. (*see* **Beckman, Daniel W.**)
Hua, Jessica, and Benjamin A. Pierce. Lethal and Sublethal Effects of Salinity on Three Common Texas Amphibians 562
Injaian, Allison (*see* **Belasen, Anat**)
Jacquemin, Stephen J., and Mark Pyron. Effects of Allometry, Sex, and River Location on Morphological Variation of Freshwater Drum *Aplodinotus grunniens* in the Wabash River, USA 740
Jelks, Howard L. (*see* **Holt, Daniel E.**)
Jerep, Fernando C., and Richard P. Vari. New Species of *Hemigrammocharax* (Characiformes: Distichodontidae) from the Northwestern Congo River Basin 31
Jordan, Frank (*see* **Holt, Daniel E.**)
Kenaley, Christopher P. (*see* **Stevenson, Duane E.**)
Killebrew, Flavius C. (*see* **Ward, Rocky**)
Kirillov, Alexandr A. (*see* **Reshetnikov, Andrey N.**)
Kluge, Arnold G. Trout. By James Owen 177
Krieger, Thomas J. (*see* **Camp, Carlos D.**)
Kuzovenko, Alexandr E. (*see* **Reshetnikov, Andrey N.**)
Kwiatkowski, Matthew A. (*see* **Sullivan, Brian K.**)
Labra, Antonieta, Gabriela Silva, Fernanda Norambuena, Nelson Velásquez, and Mario Penna. Acoustic Features of the Weeping Lizard's Distress Call 206
Lannoo, Michael J. (*see* **Williams, Perry J.**)
Lazcano, David (*see* **Borja, Miguel**)
Le, Minh D. (*see* **Nguyen, Truong Q.**)
Lehmann A., Pablo, Bruna Koech Braun, Edson H. L. Pereira, and Roberto E. Reis. A New Species of the Hypoptopomatinae Catfish *Parotocinclus* (Siluriformes: Loricariidae), from the Headwaters of the Rio Jequitinhonha, Brazil 435
Levine, Todd D. (*see* **Anderson, Thomas L.**)
Li, Kevin (*see* **Belasen, Anat**)
Lima, Albertina P. (*see* **de Fraga, Rafael**)
Lima, Flávio C. T. (*see* **Birindelli, José L. O.**)
Lima, Flávio C. T. (*see* **Teixeira, Tulio F.**)
Lind, Craig (*see* **Putman, Breanna J.**)
Lohman, Brian K., Howard I. Sirotkin, and Michael A. Bell. A Whole-Mount Method for Trypsin Clearing and Collagen Type II Antibody Staining 127
López-Fernández, Hernán (*see* **Lujan, Nathan K.**)
Lorenz, Jerome J. (*see* **Harrison, Elizabeth**)
Lovich, Jeffrey E. Turtles, Tortoises and Terrapins: A Natural History. By R. Orenstein 570
Lujan, Nathan K., Henry Agudelo-Zamora, Donald C. Taphorn, Pieter N. Booth, and Hernán López-Fernández. Description of a New, Narrowly Endemic South American Darter (Characiformes: Crenuchidae) from the Central Guiana Shield Highlands of Guyana 454
Lumbantobing, Daniel (*see* **Parenti, Lynne R.**)
Lumbantobing, Daniel N. (*see* **Tang, Kevin L.**)
MacLatchy, Deborah L. (*see* **Bosker, Thijs**)
Magnusson, William E. (*see* **de Fraga, Rafael**)
Maisey, John G., and Victor G. Springer. Chondrocranial Morphology of the Salmon Shark, *Lamna ditropis*, and the Porbeagle, *L. nasus* (Lamnidae) 378
Malabarba, Luiz R. (*see* **Villa-Verde, Leandro**)
Mandrak, Nicholas E. (*see* **Bunt, Christopher M.**)
Marks, Sharyn B. (*see* **Bondi, Cheryl A.**)
Martin, Karen L. M., Kathryn A. Hieb, and Dale A. Roberts. A Southern California Icon Surfs North: Local Ecotype of California Grunion, *Leuresthes tenuis* (Atherinopsidae), Revealed by Multiple Approaches during Temporary Habitat Expansion into San Francisco Bay 729
Martínez-Romero, Gerardo (*see* **Borja, Miguel**)
Mayden, Richard L. (*see* **Tang, Kevin L.**)
Mays, D'arcy (*see* **Sismour, Edward N.**)
McMahan, Caleb D. (*see* **Mohammadi, Shabnam**)
Melville, Jane. Tree Thinking: An Introduction to Phylogenetic Biology. By D. A. Baum and S. D. Smith 783
Mendelson, Tamra C. (*see* **Ciccotto, Patrick J.**)
Menezes, Naércio A. (*see* **Arcila, Dahiana**)
Merchant, Mark (*see* **Murray, Christopher M.**)
Miaud, Claude (*see* **Cogălniceanu, Dan**)
Miorando, Priscila S. (*see* **Pignati, Marina T.**)
Miranda, Dylria Paula da Silva (*see* **Wosiacki, Wolmar Benjamin**)
Mitchell, Joseph C., and David G. Smith. *Copeia* 1913, Number 1: Origin and Authors 189
Mitchell, Joseph C., and David G. Smith. The History of *Copeia*: Metamorphosis and the Academic Coup 361
Mitchell, Joseph C. (*see* **Smith, David G.**)
Mitchell, Joseph C. (*see* **Stewart, Margaret M.**)
Mitchem, Joshua A. (*see* **Camp, Carlos D.**)
Mohammadi, Shabnam, and Caleb D. McMahan. Venomous Reptiles of the United States, Canada, and Northern Mexico Vol. 2: *Crotalus*. By C. H. Ernst and E. M. Ernst 177
Morlett, Jesús (*see* **Borja, Miguel**)
Morris, Humberto J. (*see* **Alfonso, Yasel U.**)
Mott, Cy L. (*see* **Anderson, Thomas L.**)
Munkittrick, Kelly R. (*see* **Bosker, Thijs**)
Murray, Christopher M., Michael Easter, Mark Merchant, Amos Cooper, and Brian I. Crother. Can Reproductive Allometry Assess Population Marginality in Crocodilians? A Comparative Analysis of Gulf Coast American Alligator *(Alligator mississippiensis)* Populations 268
Muths, Erin (*see* **Swanson, Jennifer E.**)
Nacci, Diane E. (*see* **Bosker, Thijs**)
Navarrete, Ronald (*see* **Aguirre, Windsor E.**)
Nellis, Shelley C. (*see* **Sismour, Edward N.**)
Newton, Scott H. (*see* **Sismour, Edward N.**)
Nguyen, Truong Q., Trung M. Phung, Minh D. Le, Thomas Ziegler, and Wolfgang Böhme. First Record of the Genus *Oreolalax* (Anura: Megophryidae) from Vietnam with Description of a New Species 213
Nichols, John. Our Hearts Belonged to Nature 802
Norambuena, Fernanda (*see* **Labra, Antonieta**)
Nunes, Ivan, Pablo Suárez, Marcelo Gordo, and José P. Pombal, Jr. A Second Species of *Trachycephalus* Tschudi (Anura: Hylidae) with a Single Vocal Sac from the Brazilian Amazon 634
Ogle, Megan E. (*see* **Smith, Geoffrey R.**)
Oliveira, Leonardo (*see* **Dourado, Ângelo C. M.**)

Orth, Donald J. (*see* **White, Shannon L.**)
Papenfuss, Theodore J. Herpetofauna of Armenia and Nagorno-Karabakh. By M. S. Arakelyan, F. D. Danielyan, C. Corti, R. Sindaco, and A. E. Leviton 181
Parenti, Lynne R., Renny K. Hadiaty, Daniel Lumbantobing, and Fabian Herder. Two New Ricefishes of the Genus *Oryzias* (Atherinomorpha: Beloniformes: Adrianichthyidae) Augment the Endemic Freshwater Fish Fauna of Southeastern Sulawesi, Indonesia 403
Parker, Patricia G. (*see* **Ettling, Jeffery A.**)
Paull, Phillip (*see* **Doody, J. Sean**)
Peixoto, Luiz Antônio Wanderley, Guilherme Moreira Dutra, Carlos David de Santana, and Wolmar Benjamin Wosiacki. A New Species of the Electric Fish Genus *Hypopygus* (Gymnotiformes: Hypopomidae) from the Lower Amazon Basin, Brazil 232
Peixoto, Luiz Antônio Wanderley, and Wolmar B. Wosiacki. A New Species of *Characidium* (Characiformes: Crenuchidae) from the Lower Amazon 52
Peixoto, Luiz A. W. (*see* **Birindelli, José L. O.**)
Peñaloza, Claudia L., Omar Hernández, Rodolfo Espín, Larry B. Crowder, and Guillermo R. Barreto. Harvest of Endangered Sideneck River Turtles (*Podocnemis* spp.) in the Middle Orinoco, Venezuela 111
Penna, Mario (*see* **Labra, Antonieta**)
Pereira, Edson H. L. (*see* **Lehmann A., Pablo**)
Perfecto, Ivette (*see* **Belasen, Anat**)
Pezzuti, Juarez C. B. (*see* **Pignati, Marina T.**)
Phillips, Christopher A. (*see* **Allender, Matthew C.**)
Phung, Trung M. (*see* **Nguyen, Truong Q.**)
Pierce, Benjamin A. (*see* **Hua, Jessica**)
Pietsch, Theodore W., Steve W. Ross, John H. Caruso, Miles G. Saunders, and Charles R. Fisher. *In-Situ* Observations of the Deep-sea Goosefish *Sladenia shaefersi* Caruso and Bullis (Lophiiformes: Lophiidae), with Evidence of Extreme Sexual Dimorphism 660
Pietsch, Theodore W. (*see* **Vieira, Sofia**)
Pignati, Marina T., Luana F. Fernandes, Priscila S. Miorando, Paulo D. Ferreira, and Juarez C. B. Pezzuti. Effects of the Nesting Environment on Embryonic Development, Sex Ratio, and Hatching Success in *Podocnemis unifilis* (Testudines: Podocnemidae) in an Area of *Várzea* Floodplain on the Lower Amazon River in Brazil 303
Pombal, José P., Jr. (*see* **Nunes, Ivan**)
Protasova, Ekaterina N. (*see* **Reshetnikov, Andrey N.**)
Prudente, Ana L. C. (*see* **Dourado, Ângelo C. M.**)
Putman, Breanna J., Craig Lind, and Emily N. Taylor. Does Size Matter? Factors Influencing the Spatial Ecology of Northern Pacific Rattlesnakes (*Crotalus oreganus oreganus*) in Central California 485
Pyron, Mark (*see* **Jacquemin, Stephen J.**)
Radder, Rajkumar S. (*see* **Gramapurohit, Narahari P.**)
Reis, Roberto E. (*see* **Lehmann A., Pablo**)
Relyea, Rick (*see* **Hanlon, Shane M.**)
Reshetnikov, Andrey N., Sergey G. Sokolov, Igor V. Chikhlyaev, Alexandr I. Fayzulin, Alexandr A. Kirillov, Alexandr E. Kuzovenko, Ekaterina N. Protasova, and Maxim O. Skomorokhov. Direct and Indirect Interactions between an Invasive Alien Fish (*Perccottus glenii*) and Two Native Semi-aquatic Snakes 103
Richards-Dimitrie, Teal, Susan E. Gresens, Scott A. Smith, and Richard A. Seigel. Diet of Northern Map Turtles (*Graptemys geographica*): Sexual Differences and Potential Impacts of an Altered River System 477
Robb, Joseph R. (*see* **Williams, Perry J.**)
Roberts, Dale A. (*see* **Martin, Karen L. M.**)
Robson, Laura E., and Gabriel Blouin-Demers. Eastern Hognose Snakes (*Heterodon platirhinos*) Avoid Crossing Paved Roads, but Not Unpaved Roads 507
Rocha, Luiz A. Reef Fishes of the East Indies. Volumes I–III. By Gerald R. Allen and Mark V. Erdmann 567
Rodríguez-Schettino, Lourdes (*see* **Alfonso, Yasel U.**)
Rogers, Lock, Kelly Smith, and Megan Williams. Sex-Specific Planktonic Larval Durations in the Bluehead Wrasse, *Thalassoma bifasciatum* 493
Roşioru, Daniela (*see* **Cogălniceanu, Dan**)
Ross, Steve W. (*see* **Pietsch, Theodore W.**)
Ryerson, William G. Jumping in the Salamander *Desmognathus ocoee* 512
Salerno, Patricia E. Color Catalogue for Field Biologists. By Gunther Köhler 569
Sanaiotti, Tania (*see* **de Fraga, Rafael**)
Sánchez, Elda (*see* **Borja, Miguel**)
Saunders, Miles G. (*see* **Pietsch, Theodore W.**)
Savage, Jay M., Cristina A. Ugarte, and Maureen A. Donnelly. A New Species of Earless Toad (Bufonidae: *Incilius*) from Western Panama 8
Scott, John W. (*see* **Allender, Matthew C.**)
Seegert, Greg. Fishes of Indiana: A Field Guide. By Thomas P. Simon, Illustrations by Joseph R. Tomelleri 352
Seigel, Richard A. (*see* **Richards-Dimitrie, Teal**)
Sessions, Stanley K. The Chromosomes of Terraranan Frogs: Insights into Vertebrate Cytogenetics. By M. Schmid, J. P. Bogart, and S. B. Hedges (eds.) 179
Shervette, Virginia R. (*see* **Aguirre, Windsor E.**)
Shoemaker, Kevin T., and James P. Gibbs. Genetic Connectivity among Populations of the Threatened Bog Turtle (*Glyptemys muhlenbergii*) and the Need for a Regional Approach to Turtle Conservation 324
Silva, Gabriela (*see* **Labra, Antonieta**)
Simmons, Jeffrey W. Chronology of the Invasion of the Tennessee and Cumberland River Systems by the Mississippi Silverside, *Menidia audens*, with Analysis of the Subsequent Decline of the Brook Silverside, *Labidesthes sicculus* 292
Sirotkin, Howard I. (*see* **Lohman, Brian K.**)
Sismour, Edward N., Shelley C. Nellis, Scott H. Newton, D'arcy Mays, and Michael L. Fine. An Experimental Study of Consumption of Channel Catfish *Ictalurus punctatus* by Largemouth Bass *Micropterus salmoides* when Alternative Prey Are Available 277
Skomorokhov, Maxim O. (*see* **Reshetnikov, Andrey N.**)
Smith, David G., and Joseph C. Mitchell. 100th Anniversary of *Copeia* 1
Smith, David G. (*see* **Mitchell, Joseph C.**)
Smith, Geoffrey R., Christopher J. Dibble, Andrew J. Terlecky, Christopher B. Dayer, Allison B. Burner, and Megan E. Ogle. Effects of Invasive Western Mosquitofish and Ammonium Nitrate on Green Frog Tadpoles 248
Smith, Kelly (*see* **Rogers, Lock**)
Smith, Scott A. (*see* **Richards-Dimitrie, Teal**)
Sokolov, Sergey G. (*see* **Reshetnikov, Andrey N.**)
Spotila, James R. Sea Turtles of the Eastern Pacific, Advances in Research and Conservation. By Jeffrey A. Seminoff and Bryan P. Wallace (eds.) 568

Springer, Victor G. (*see* **Maisey, John G.**)
Steffen, John E. (*see* **Alfonso, Yasel U.**)
Stevenson, Duane E., and Christopher P. Kenaley. Revision of the Manefish Genera *Caristius* and *Platyberyx* (Teleostei: Percomorpha: Caristiidae), with Descriptions of Five New Species 415
Stewart, Donald J. Re-description of *Arapaima agassizii* (Valenciennes), a Rare Fish from Brazil (Osteoglossomorpha: Osteoglossidae) 38
Stewart, Donald J. A New Species of *Arapaima* (Osteoglossomorpha: Osteoglossidae) from the Solimões River, Amazonas State, Brazil 470
Stewart, Margaret M., and Joseph C. Mitchell. Herndon Glenn Dowling, Jr. 166
Stuart-Fox, Devi. Concealing Coloration in Animals. By J. Diamond and A. B. Bond 782
Suárez, Pablo (*see* **Nunes, Ivan**)
Sullivan, Brian K., James M. Walker, Harry L. Taylor, James E. Cordes, Matthew A. Kwiatkowski, Keith O. Sullivan, Justin R. Sullivan, Marlis R. Douglas, and Michael E. Douglas. Morphological Diagnosability of *Aspidoscelis arizonae* (Squamata: Teiidae) as an Indication of Evolutionary Divergence in the *Aspidoscelis inornata* Complex 366
Sullivan, Justin R. (*see* **Sullivan, Brian K.**)
Sullivan, Keith O. (*see* **Sullivan, Brian K.**)
Swanson, Jennifer E., Larissa L. Bailey, Erin Muths, and W. Chris Funk. Factors Influencing Survival and Mark Retention in Postmetamorphic Boreal Chorus Frogs 670
Sweet, Samuel S. Field Guide to Amphibians and Reptiles of California. Revised edition. By R. C. Stebbins and S. M. McGinnis 785
Székely, Diana (*see* **Cogălniceanu, Dan**)
Székely, Paul (*see* **Cogălniceanu, Dan**)
Tang, Kevin L., Daniel N. Lumbantobing, and Richard L. Mayden. The Phylogenetic Placement of *Oxygaster* van Hasselt, 1823 (Teleostei: Cypriniformes: Cyprinidae) and the Taxonomic Status of the Family-Group Name Oxygastrinae Bleeker, 1860 13
Taphorn, Donald C. (*see* **Lujan, Nathan K.**)
Taylor, Emily N. (*see* **Putman, Breanna J.**)
Taylor, Harry L. (*see* **Sullivan, Brian K.**)
Teixeira, Tulio F., Flávio C. T. Lima, and Jansen Zuanon. A New *Hyphessobrycon* Durbin from the Rio Teles Pires, Rio Tapajós Basin, Mato Grosso State, Brazil (Characiformes: Characidae) 612
Terlecky, Andrew J. (*see* **Smith, Geoffrey R.**)
Terrell, Vanessa C. K. (*see* **Williams, Perry J.**)
Trexler, Joel C. (*see* **Harrison, Elizabeth**)
Ugarte, Cristina A. (*see* **Savage, Jay M.**)
Vari, Richard P., and Carl J. Ferraris, Jr. Two New Species of the Catfish Genus *Tatia* (Siluriformes: Auchenipteridae) from the Guiana Shield and a Reevaluation of the Limits of the Genus 396
Vari, Richard P. (*see* **Arcila, Dahiana**)
Vari, Richard P. (*see* **Ferraris, Carl J., Jr.**)
Vari, Richard P. (*see* **Jerep, Fernando C.**)
Velásquez, Nelson (*see* **Labra, Antonieta**)
Vieira, Sofia, Manuel Biscoito, Helena Encarnação, João Delgado, and Theodore W. Pietsch. Sexual Parasitism in the Deep-sea Ceratioid Anglerfish *Centrophryne spinulosa* Regan and Trewavas (Lophiiformes: Centrophrynidae) 666
Vilano, Ma. de Lourdes Lozano (*see* **Echelle, Anthony A.**)
Villa-Verde, Leandro, Juliano Ferrer, and Luiz R. Malabarba. A New Species of *Listrura* from Laguna dos Patos System, Brazil: The Southernmost Record of the Glanapteryginae (Siluriformes: Trichomycteridae) 641
Vitt, Laurie J. Geckos: The Animal Answer Guide. By A. M. Bauer 781
Wake, Marvalee H. (*see* **Donnelly, Maureen A.**)
Walker, James M. (*see* **Sullivan, Brian K.**)
Walls, Susan C. (*see* **Brown, Mary E.**)
Ward, Rocky, Joel B. Babitzke, and Flavius C. Killebrew. Genetic Population Structure of Cagle's Map Turtle (*Graptemys caglei*) in the Guadalupe and San Marcos Rivers of Texas—A Landscape Perspective 723
Watkins-Colwell, Gregory J. Contributions to the History of Herpetology. Volume 3. By K. Adler (ed.) 351
Wegner, Nicholas C. (*see* **Baird, Troy A.**)
White, Shannon L., and Donald J. Orth. Ontogenetic and Comparative Morphology of Clinch Dace (*Chrosomus* sp. cf. *saylori*) 750
Whiteman, Howard H. (*see* **Anderson, Thomas L.**)
Williams, Megan (*see* **Rogers, Lock**)
Williams, Perry J., Nathan J. Engbrecht, Joseph R. Robb, Vanessa C. K. Terrell, and Michael J. Lannoo. Surveying a Threatened Amphibian Species through a Narrow Detection Window 552
Wilson, Wade D. (*see* **Echelle, Anthony A.**)
Wooten, Jessica A. (*see* **Camp, Carlos D.**)
Wosiacki, Wolmar Benjamin, and Dylria Paula da Silva Miranda. Description of a New Small Species of the Genus *Cyphocharax* (Characiformes: Curimatidae) from the Lower Amazon Basin 627
Wosiacki, Wolmar B. (*see* **Birindelli, José L. O.**)
Wosiacki, Wolmar B. (*see* **Peixoto, Luiz Antônio Wanderley**)
Wosiacki, Wolmar Benjamin (*see* **Peixoto, Luiz Antônio Wanderley**)
Wylie, Daniel B. (*see* **Allender, Matthew C.**)
Wylie, Sarah J. (*see* **Allender, Matthew C.**)
Ziegler, Thomas (*see* **Nguyen, Truong Q.**)
Zuanon, Jansen (*see* **de Pinna, Mário C. C.**)
Zuanon, Jansen (*see* **Teixeira, Tulio F.**)
Zugasti-Cruz, Alejandro (*see* **Borja, Miguel**)